



# International Journal of Medical Anesthesiology

E-ISSN: 2664-3774  
P-ISSN: 2664-3766  
IJMA 2019; 2(2): 256-260  
Received: 22-05-2019  
Accepted: 25-06-2019

**Dr. Karri Bala Krishna**  
Assistant Professor of  
Anaesthesia Great Eastern  
Medical School & Hospital,  
Ragolu, Srikakulam, Andhra  
Pradesh, India

**Dr. Nellimarala Appala Raju**  
Assistant Professor of  
Anaesthesia Great Eastern  
Medical School & Hospital,  
Ragolu, Srikakulam,  
Andhra Pradesh, India

**P. Kalyan Chakravarthy**  
Associate professor Great  
Eastern Medical School &  
Hospital, Ragolu, Srikakulam,  
Andhra Pradesh, India

**Corresponding Author:**  
**Dr. Nellimarala Appala Raju**  
Assistant Professor of  
Anaesthesia Great Eastern  
Medical School & Hospital,  
Ragolu, Srikakulam,  
Andhra Pradesh, India

## Pre-operative anxiety: A prevalence study in Northern Andhra Pradesh

**Dr. Karri Bala Krishna, Dr. Nellimarala Appala Raju and P. Kalyan Chakravarthy**

### Abstract

The prospect of undergoing surgery under anaesthesia can induce significant anxiety in patients. Excess anxiety can lead to several deleterious effects like perioperative cardiac events, increased anaesthetic requirements, higher postoperative pain scores, and prolonged hospital stay. Various factors can influence anxiety levels during surgery. The aim of the study was to estimate the prevalence of anxiety in adult patients scheduled for surgery in a suburban teaching institute in India where the majority of patients belong to lower socioeconomic strata.

**Methods:** One hundred twenty consecutive patients scheduled for various elective surgeries were asked 6 questions according to the Amsterdam preoperative anxiety and information scale to assess their anxiety levels just before shifting into the Operating room. Probable contributing factors noted were age, sex, socioeconomic status, type of surgery, duration of hospital stay, and history of previous surgery. Any specific factors causing anxiety like failure or complications of the surgery, recovery from anaesthesia, needle pricks were also asked with leading questions.

**Results:** The overall prevalence of anxiety was 36.3%. The prevalence in male patients and female patients was 32.4% and 40.2% respectively. There was a moderate correlation between the anxiety scores and the need to know scores. The effects of sex, previous surgery, the length of hospital stay were not significant upon the anxiety scores.

**Conclusion:** The overall prevalence of anxiety in the southern part of India was lower when compared to many of the reported studies; however, the prevalence was higher among the female patients.

**Keywords:** pre-operative anxiety, Northern Andhra Pradesh, anaesthetic

### Introduction

Anxiety is defined as "A the feeling of worry, nervousness, or unease about something with an uncertain outcome". The incidence of preoperative anxiety can be as high as 80%.<sup>[1]</sup> Excessive anxiety has deleterious effects like hypertension and tachycardia due to sympathetic stimulation and can also increase anaesthetic requirements<sup>[2]</sup> and Create complications like prolonged recovery, greater postoperative pain, etc.<sup>[3]</sup> the neurohumoral response to anxiety causes elevated intraoperative cortisol levels.<sup>[4]</sup> Anxiety can be an independent risk factor that predicts postoperative cardiac events.<sup>[5]</sup> Patient-related factors like age, sex, the level of education and economic status, pain tolerance, previous surgery and anaesthesia exposure, existing psychiatric problems, social security can affect the anxiety levels. Procedure-related factors like major surgeries, chronicity of the problem, emergency and unscheduled procedures, and destructive procedures might play a role. The attitude of the health care providers, communication skills of the doctors, the overall impression of the institution, beliefs, and opinions of neighboring patients also exert influence. Preoperative Anaesthetic Information, Anaesthetic Catastrophizing, and Imminence of Surgery were shown to be statistically significantly associated with an overall increased level of anxiety.<sup>[6]</sup> Other factors like fear of disability, medical errors, noise levels, unpleasant smell, financial loss, cancellations also play a role.

Pharmacological anxiolysis is a commonly adopted strategy to allay anxiety, the effectiveness of which is variable. Among the premedication drugs used for anxiolysis, benzodiazepines have been shown to provide consistent anxiolysis. Diazepam is one of the most common benzodiazepines used for this purpose.<sup>[7]</sup> Anxiolytic effects are seen at low doses because of diazepam's interaction with BZ2 receptors in the limbic system.<sup>8</sup> However, the value of routine anxiolytic premedication is questioned in the era of better patient education.<sup>9</sup> An empathic attitude by the anaesthesiologist during the preoperative visit can decrease anxiety when continuous care is provided by the same anaesthesiologist (One

patient one anaesthesiologist).<sup>[10]</sup> this may not be possible in countries like India where the patient load is high. Preoperative multimedia information about the procedure can also decrease anxiety immediately before the surgery.<sup>[11]</sup> Well-informed patients are less anxious.<sup>12</sup>Preoperative anxiety among adults is an under-reported problem in India and only a few studies have addressed the issue.<sup>[13]</sup> The Amsterdam preoperative anxiety and Information scale is a validated tool for the measurement of anxiety. The scale has been translated into various languages like German<sup>14</sup>, Mexican<sup>[15]</sup>, Thai<sup>[16]</sup>, Korean<sup>[17]</sup>, Japanese<sup>[18]</sup>, and Srilankan.<sup>19</sup> We have used an Indian language (Telugu) version of APAIS as a measurement tool for anxiety. The aim of this study was to estimate the prevalence of anxiety in adult general surgical premedicated patients just before shifting into the operating room.

**Methods**

After ethics committee approval, one hundred twenty consecutive patients undergoing elective surgeries were recruited as a convenient sample for this observational cross-sectional study. The period of the study was from July to September 2018. Consent was taken just before the questionnaire was answered.

Inclusion criteria were all adult patients undergoing elective surgery, had received premedication the previous night, willing to participate in the study, and able to answer the questions. Patients with pre-existing psychiatric illnesses, those who did not receive premedication, those having difficulty understanding and communicating, or unwilling to participate were excluded.

The primary objective of this study was to estimate the prevalence of anxiety in adult general surgical premeditated patients just before shifting into the operating room. Secondary objectives include assessing the correlation of sex, socio-economic status, previous surgery, and length of hospital stay upon anxiety.

In the preoperative holding area, the questions from the APAIS questionnaire were asked by a single anaesthesia resident to assess the degree of anxiety. The patients were comfortably seated or lying down position. Privacy and confidentiality were ensured before asking questions.

The APAIS consists of 6 questions, 2 each for anaesthesia and surgery-related anxiety, and 2 more for information desired component related to anaesthesia and surgery. Each question is scored on a 1 to 5 Likert scale, 1 being minimal and 5 denoting extreme values.

**Table 1: APAIS Questionnaire**

	Not at all 1 2 3 4 5 extremely
1. I am worried about the anaesthetic	
2. The anaesthetic is on my mind continually	
3. I would like to know as much as possible about the anaesthetic	
4. I am worried about the procedure	
5. The procedure is on my mind continually	
6. I would like to know as much as possible about the procedure	
The subscales	
Anesthesia-related anxiety	Sum A = 1 + 2
Surgery-related anxiety	Sum S = 4 + 5
Information desired component	Sum I= 3 + 6
Combined anxiety component	Sum C = sum A + sum S ( 1 + 2 + 4 +5)

Other data collected were:

1. Age, sex, nature of surgery
2. Socio-economic status. (Modified Kuppuswami socioeconomic status scale 2012)
3. Length of hospital stay
4. Previous surgery
5. Any Specific anxiety-inducing factor (What do you worry the most regarding surgery?)

APAIS anxiety scores of more than 10 were considered as anxiety. Prevalence of anxiety is presented as the percentage. Pearson’s correlation was used to correlate anxiety score and desire for information score. Chi-Squared test was used to establish the effect of sex, previous surgery and length of hospital stay (Short < 1 week and long > 1 week). Kruskal-Wallis test was used to compare anxiety scores with respect to socio-economic status. An alpha error of 5% was allowed.

**Results**

The overall prevalence of anxiety (APAIS anxiety score >10) was 36.3%. The prevalence of male patients and female patients was 32.4% and 40.2% respectively. (Table 2) The higher prevalence in female patients was statistically significant. The effect of social status upon anxiety scores

was not significant as analyzed by the Kruskal-Wallis test. The majority of the patients fell into social status III and IV. Chi-square test showed no significant influence of previous surgery and length of hospital stay (short-17 days or less, long- more than 7 days) upon anxiety scores.

The mean anaesthesia, surgery, and total anxiety scores were 3.56, 3.81, and 7.27 respectively. There was a moderate correlation between the need to know scores with anaesthesia anxiety score and total anxiety sco

**Table 2: Factors affecting perioperative anxiety**

Parameter		APAIS >10 (%)
Sex	Male (n=62)	32.4%
	Female (n=58)	40.2%
Socio economic status (SES)	SES 4 or less (n=108)	36.8%
	SES 5 and above (n=12)	42%
Duration of hospital stay before surgery	7 days or less (n=65)	38.7%
	8 days or more (n=55)	24.8%
Previous surgery	Yes (n=84)	34.3%
	No (n=36)	28%

## Discussion

Patient anxiety is often neglected during anaesthesia and surgery, and the effects are deleterious. Anxiety may develop several days before the surgery and might continue to the post-operative period. Patients with a higher level of preoperative anxiety have higher postoperative anxiety scores. The time of anxiety assessment is also important. Several studies assess anxiety on the day before surgery. Patients can have maximum anxiety just before anaesthesia induction.<sup>[21]</sup> Patients may have higher levels of anxiety at the pre-operative holding area as compared to the ward or the operating room.<sup>[22]</sup>

Regarding tools for measurement of anxiety, the STAI (State-trait anxiety Inventory Questionnaire) is considered a gold standard. It consists of 20 statements and each statement can take values from not at all, somewhat, moderately, or very much so. Even though STAI is available in several languages, the statements become difficult to translate and become incomprehensible to patients with low educational levels. For example, "I feel tense;" and "I feel jittery" might mean the same in translation, and a more literary word loses its meaning to the layman. Hence the APAIS scoring which is simple, easy to comprehend, and correlates well with other tools of measurement of anxiety was used. Other commonly used scales are Hospital Anxiety and depression score and visual analog anxiety scale. The HADS scale is not specific for preoperative anxiety and the VAS scale does not distinguish surgical and anaesthesia-related anxiety. The APAIS score was specially designed for anxiety assessment by anaesthesiologists and has a high degree of correlation with STAI scores.<sup>[23]</sup> APAIS anxiety scores of more than 10 were considered as significant anxiety. This level was found to have an acceptable balance between sensitivity and specificity.<sup>[14]</sup>

The prevalence of anxiety in preoperative patients in various studies ranges from 51% to 76.7%.<sup>[19]</sup> A range of 60% to 80% has been suggested.<sup>[11]</sup> In a study with 79 patients in India, the prevalence of pre-operative anxiety was 45.7%.<sup>[12]</sup> But in our study, the prevalence was 31% which is lower than these reported studies.

Anxiolytic premedication the night before surgery is a common practice, and in our study, most patients had received oral diazepam. This is one of the reasons the prevalence of anxiety is low in our study. The prevalence of anxiety was higher in females in agreement with previous studies.<sup>[1]</sup><sup>[23]</sup> The correlation between total anxiety scores and need for information score was moderate similar to other studies.<sup>[23]</sup>

Most of the patients were of lower and upper-lower socioeconomic status. Economic and social status plays an important role in anxiety. Income levels have been shown to be an independent predictor of pre-operative anxiety. For a unit increase in income, preoperative state anxiety score increases by 0.002.<sup>[25]</sup> This might also explain the lower prevalence of anxiety in our study group. Few reported studies have taken a sample size ranging from 51 to 2002,24, our sample size was based on a previous study conducted in Srilankan population<sup>19</sup> and is a sample of convenience.

The factor that induces most anxiety varies across countries and reflects the cultural and traditional values of society. Whereas in the west, waiting, pain and awareness, loss of control, etc are anxiety-inducing factors<sup>1</sup>, concerns about family, financial losses were shown to be the important

factors in a study from Pakistan<sup>[26]</sup>, and fear of death and unknown was the major factor in an Ethiopian study<sup>[25]</sup>, fear of postponement and mishaps were the leading cause of anxiety in a Nigerian study.<sup>[27]</sup> No specific anxiety-inducing factor emerged in our study.

What we have measured is the prevalence of anxiety among a set of surgical patients, as we have not assessed whether the patients were already anxious (Prone to anxiety -high trait anxiety) or the anxiety is entirely due to the surgery and anaesthesia (State anxiety). The APAIS score does not distinguish state anxiety (anxiety during a threat) and Trait anxiety (tendency to respond with state anxiety) in individuals, unlike the STAI questionnaire which has 20 questions each for state and trait anxiety, separately. However, the APAIS correlates well with the total STAI scores. If we had taken a set number of patients without pre-existing anxiety and followed them up to the perioperative period (or a set time, like one year) and then measured how many of them develop anxiety then that would indicate the Incidence of anxiety. What we have measured is a point prevalence of anxiety across a cross-section of the population during the preoperative period. Prevalence does not distinguish "Old" cases and "New" cases, that is those who were already anxious and those who experience anxiety only before the surgery. Strengths and Limitations We have employed a regional language version of the APAIS score to assess preoperative anxiety in premedicated patients just before shifting inside the operating room, in contrast to most studies that assessed anxiety the day before surgery or without premedication. Our study design resembles the real-world scenario more closely, enhancing the external validity and applicability of the results.

The regional language version of the primary measurement tool is not validated, a major limitation of our study. The sample size of hundred was selected as it was more practical, might not be sufficient to extrapolate the results to a wider population. Also, most of the patients were from low socio-economic status and the results might not apply to other population categories. The estimated prevalence was lower as compared to other studies. The reason for low anxiety can be multifactorial, and a larger sample with multivariate regression analysis might delineate the influence of individual factors upon anxiety.

In conclusion, we have evaluated pre-operative anxiety in premedicated patients using a regional language Tamil version of the APAIS scale just before surgery. The result shows a prevalence of 31% and is significantly more in female patients when compared to male patients. The anxiety scores correlate with the need for information scores and dissemination of more procedure-related information may help in reducing the anxiety.

## References

1. Mitchell M. Anaesthesia type, gender and anxiety. *J Perioper Pract* 2013; 23(3):41-7. [PMid:23586187]
2. Maranets I, Kain ZN. Preoperative anxiety and intraoperative anaesthetic requirements. *Anesth Analg* 1999; 89(6):1346-51. [PMid:10589606]
3. IP HYV, Abrishami A, Peng PWH, Wong J, Chung F. Predictors of postoperative pain and analgesic consumption: A qualitative systematic review. *Anesthesiology* 2009; 111(3):657-77. <https://doi.org/10.1097/ALN.0b013e3181aae87a> [PMid:19672167]

4. Pearson S, Maddern GJ, Fitridge R. The role of pre-operative state-anxiety in the determination of intra-operative neuroendocrine responses and recovery. *Br J Health Psychol* 2005; 10:299-310.  
<https://doi.org/10.1348/135910705X26957>  
[PMid:15969856]
5. Szekely A, Balog P, Benko E, Breuer T, Szekely J, Kertai MD, *et al.* Anxiety predicts mortality and morbidity after coronary artery and valve surgery- a 4-year follow-up study. *Psychosom Med* 2007; 69(7):625-31.  
<https://doi.org/10.1097/PSY.0b013e31814b8c0f>  
[PMid:17724254]
6. Mitchell M. General anaesthesia and day-case patient anxiety. *J Adv Nurs* 2010; 66(5):1059-71.  
<https://doi.org/10.1111/j.1365-2648.2010.05266.x>  
[PMid:20337788]
7. Fox C, Liu H, Kaye AD, Manchikanti L, Trescot AM, Christo PJ. Clinical aspects of pain medicine and interventional pain management: a comprehensive review. Paducah, KY: ASIP Publishing. Antianxiety agents. 2011:543-52.
8. Griffin CE, Kaye AM, Bueno FR, Kaye AD. Benzodiazepine pharmacology and central nervous system mediated effects. *Ochsner J.* 2013; 13(2):214-23. [PMid:23789008] [PMCID:PMC3684331]
9. Beydon L, Rouxel A, Camut N, Schinkel N, Malinovsky J-M, Aveline C, *et al.* Sedative premedication before surgery-A multicentre randomized study versus placebo. *Anaesth Crit Care Pain Med.* 2015; 34(3):165-71.  
<https://doi.org/10.1016/j.accpm.2015.01.005>  
[PMid:26004882]
10. Soltner C, Giquello JA, Monrigal-Martin C, Beydon L. Continuous care and empathic anaesthesiologist attitude in the preoperative period: impact on patient anxiety and satisfaction. *Br J Anaesth* 2011; 106(5):680-6.  
<https://doi.org/10.1093/bja/aer034> [PMid: 21414978]
11. Jjala HA, French JL, Foxall GL, Hardman JG, Bedford NM. Effect of preoperative multimedia information on perioperative anxiety in patients undergoing procedures under regional anaesthesia. *Br J Anaesth* 2010; 104(3):369-74.  
<https://doi.org/10.1093/bja/aeq002> [PMid:20124283]
12. Nikumb VB, Banerjee A, Kaur G, Chaudhury S. Impact of doctor-patient communication on preoperative anxiety: Study at industrial township, Pimpri, Pune. *Ind Psychiatry J* 2009; 18(1):19-21.  
<https://doi.org/10.4103/0972-6748.57852>  
[PMid:21234156] [PMCID:PMC3016692]
13. Akkamahadevi P, Subramanian VV. The efficacy of different methods of pre-operative counselling on perioperative anxiety in patients undergoing regional anaesthesia. *Indian J Anaesth* 2016; 60(1):58.  
<https://doi.org/10.4103/0019-5049.174812>[PMid:26962258] [PMCID:PMC4782426]
14. Berth H, Petrowski K, Balck F. The Amsterdam Preoperative Anxiety and Information Scale (APAIS) - the first trial of a German version. *Psycho-Soc Med* 2007; 4:Doc01.  
[PMid:19742298] [PMCID:PMC2736533]
15. Del Carmen Gavito M, Corona MA, Villagran ME, Morales J, Tellez JL, Ortega-Soto HA. La informacion anestésica quirúrgica: su efecto sobre la ansiedad y el dolor de los pacientes toracotomizados. *Rev Inst Nal Enf Resp Mex.* 2000; 13(3):153-6.
16. Kunthonluxamee A, Pitimana-aree S, Laurujisawat P. Validity and Reliability of the Amsterdam Preoperative Anxiety and Information Scale (APAIS); Thai version in adult Thai pre-operative patients. *J Psychiatr Assoc Thailand.* 2013; 54(1):83-92.
17. Shin WJ, Kim YC, Yeom JH, Cho SY, Lee DH, Kim DW. The Validity of Amsterdam Preoperative Anxiety Information Scale in the Assessment of the Preoperative Anxiety - Compared with hospital anxiety depression scale and visual analogue scale. *Korean J Anesthesiol.* 1999; 37(2):179-87.  
<https://doi.org/10.4097/kjae.1999.37.2.179>
18. Nishimori M, Moerman N, Fukuhara S, van Dam FS, Muller MJ, Hanaoka K, *et al.* Translation and validation of the Amsterdam preoperative anxiety and information scale (APAIS) for use in Japan. *Qual Life Res* 2002; 11(4):361-4.  
<https://doi.org/10.1023/A:1015561129899>  
[PMid:12086121]
19. Matthias AT, Samarasekera DN. Preoperative anxiety in surgical patients-experience of a single unit. *Acta Anaesthesiol Taiwanica.* 2012; 50(1):3-6.  
<https://doi.org/10.1016/j.aat.2012.02.004>  
[PMid:22500906]
20. Dudala SR. Updated Kuppuswamy's socioeconomic scale for 2012. *J NTR Univ Health Sci* 2013; 2:201-2.  
<https://doi.org/10.4103/2277-8632.117195>
21. Haugen AS, Eide GE, Olsen MV, Haukeland B, Remme AR, Wahl AK. Anxiety in the operating theatre: a study of frequency and environmental impact in patients having local, plexus or regional anaesthesia. *J Clin Nurs* 2009; 18(16):2301-10.  
<https://doi.org/10.1111/j.1365-2702.2009.02792.x>  
[PMid:19583663]
22. Pokharel K, Bhattarai B, Tripathi M, Khatiwada S, Subedi A. Nepalese patients anxiety and concerns before surgery. *J Clin Anesth* 2011; 23(5):372-8.  
<https://doi.org/10.1016/j.jclinane.2010.12.011>  
[PMid:21802628]
23. Moerman N, van Dam FS, Muller MJ, Oosting H. The Amsterdam Preoperative Anxiety and Information Scale (APAIS). *Anesth Analg.* 1996; 82(3):445-51. [PMid:8623940]
24. Akinsulore A, Owojuyigbe AM, Faponle AF, Fatoye FO. Assessment of preoperative and postoperative anxiety among elective major surgery patients in a tertiary hospital in Nigeria. *Middle East J Anaesthesiol* 2015; 23(2):235-40. [PMid:26442401]
25. Nigussie S, Belachew T, Wolancho W. Predictors of preoperative anxiety among surgical patients in Jimma University Specialized Teaching Hospital, South Western Ethiopia. *BMC Surg* 2014; 14:67.  
<https://doi.org/10.1186/1471-2482-14-67>  
[PMid:25189274] [PMCID:PMC4167307]
26. Jafar MF, Khan FA. Frequency of preoperative anxiety in Pakistani surgical patients. *J Pak Med Assoc* 2009; 59(6):359-63. [PMid:19534368]
27. Ebirim LN, Tobin M. Factors Responsible For Pre-Operative Anxiety In Elective Surgical Patients At A University Teaching Hospital: A Pilot Study. *Internet J*

Anesthesiol [Internet]. 2010 Dec 31 [cited 2017 Jan 15];29(2). Available from:  
<http://ispub.com/IJA/29/2/8534>