



International Journal of Medical Anesthesiology

E-ISSN: 2664-3774
P-ISSN: 2664-3766
www.anesthesiologypaper.com
IJMA 2020; 3(2): 24-28
Received: 22-02-2020
Accepted: 06-03-2020

Dr. Rajeev Nair
MD Anesthesia, HOD
Department of
Anaesthesiology and critical
care Command hospital Air
force, Bangalore, Karnataka,
India.

Dr. Satish Kumar Mishra
MD, DM Cardiac Anesthesia,
Department of
Anaesthesiology and critical
care Command hospital Air
force, Bangalore, Karnataka,
India.

Dr. Gobind Ji Thakur
Mch CTVS, Army Hospital R
& R Delhi department of
CTVS, Delhi, India.

Dr. Shalendra Singh
MD Anesthesia, Department of
Anaesthesiology and critical
care, Armed forces medical
College, Pune, Maharashtra,
India.

Corresponding Author:
Dr. Satish Kumar Mishra
MD, DM Cardiac Anesthesia,
Department of
Anaesthesiology and critical
care Command hospital Air
force, Bangalore, Karnataka,
India

A comparative study between International normalized Ratio (INR) guided versus clinical discretion guided blood component replacement therapy in patients undergoing major surgery

Dr. Rajeev Nair, Dr. Satish Kumar Mishra, Dr. Gobind Ji Thakur and Dr. Shalendra Singh

DOI: <https://doi.org/10.33545/26643766.2020.v3.i2a.120>

Abstract

Bleeding is a frequent complication during surgery. The intraoperative administration of blood products, including packed red blood cells, platelets and fresh frozen plasma (FFP), is often lifesaving. Complications of blood transfusions contribute considerably to perioperative costs and blood product resources are limited. The aim of the present study was to evaluate and compare the usage of INR guided vs clinician discretion based component replacement therapy which will optimize the use of FFP and may even result in less blood loss during surgery. Materials and Method: This study was conducted in surgical patients in a large tertiary care centre. Ethical clearance was taken from the local ethics committee at the proposal stage itself. Patients consent was taken after providing all necessary information prior to surgery. This study was conducted on 200 patients. The groups were randomised to two groups of Gp 1- study (S) -100 patients (point of care based transfusion management) Gp 2- control (C)- 100 patients (physician discretion based management). After premedication with intravenous morphine (0.05- 0.1 mg/kg body weight), Glycopyrrolate and ondansetron, General anaesthesia was induced with thiopentone sodium. Endotracheal intubation achieved after vecuronium. Anaesthesia was maintained using low flow nitrous oxide: oxygen mixture (fresh gas flow of 1ltr each) and Isoflurane (1 MAC) via a closed circuit cycle absorber system and mechanical ventilation with 5 – 10 ml/kg. Standard monitoring will include HR, ECG (two lead), blood pressure (NIBP/ IABP), SpO₂, Naso pharyngeal temperature. Patient warming was done with warm air blower (with a target temp above 36⁰C). In the control group physician discretion was used to guide Blood platelet or FFP transfusion based on the institutional protocol. These are blood if Hb <8 gm %, platelets if <50,000 or between 50,000-80,000 with ongoing blood loss and FFP if bleeding >20% or >2 units blood given and repeated if physician desires. However these are guidelines and physician discretion based on clinical judgment is freely allowed. In the test group if bleeding is estimated to be more than 205 CBC and POC INR is done. If Hb < 8 gm % blood is transfused, if platelets <50,000 then 1 Single Donor Platelets or 6 Random Donor Platelets are transfused. 2 units of FFP are transfused if POC INR >1.8. Test was repeated after 30mins and 2 more units of FFP was given if INR >1.5. Result: there was no difference in age distribution sex weight of patients between the two group, however the need for blood component therapy was in higher in patients who were treated based on clinical discretion compared to INR guided treatment. The mean duration of post-operative ventilation required for the patient to be extubated with stable haemodynamic parameters in the test group was 48.55 hrs compared to 78.4 hrs for the control group. The maximum duration of post-operative ventilation in test group was 212 hrs, compared to 281 hrs for the control group. While the minimum duration was comparable being 18 hrs for test group and 19 hrs for control group. One-Sample Kolmogorov-Smirnov Test was done in each group to ensure normal distribution due to wide distribution of data. From our study we concluded that when INR guided blood component therapy was done the need for blood products was much lesser and the need for post-operative mechanical ventilation was much lesser in inr guided group compared to clinical discretion based therapy.

Keywords: INR, International Normalized Ratio, FFP: Fresh frozen plasma, POC: Point of care

Introduction

Karl Landsteiner's discovery of ABO blood groups in 1900 began the modern era of transfusion medicine. In 1915 Richard Lewisohn introduced anticoagulation with sodium citrate. Blood transfusion was extensively used in World War I and II, with a dictum of

“When in doubt transfuse”. Bleeding is a frequent complication during surgery. The intraoperative administration of blood products, including packed red blood cells, platelets and fresh frozen plasma (FFP), is often life saving. Complications of blood transfusions contribute considerably to perioperative costs and blood product resources are limited. With the emergence of HIV, Hepatitis and other blood borne illnesses has made a serious rethink on blood transfusion. Consequently, strategies to optimize the decision to transfuse are needed.

Variation in blood component use among individuals and among hospitals suggests that current transfusion practice is not ideal. Blood transfusion practice appears to be based more on local custom than on evidence. Evidence suggests that some blood transfusion practices may be harmful. There is worldwide concern over inappropriate use of blood components in view of their costs and numerous complications which include immunosuppression, all immunization and risks of transfusion – transmitted infection, in addition to hemolytic and non hemolytic reactions.

Bleeding during surgery is a dynamic process and may result in major blood loss and coagulopathy due to dilution and consumption. The indication for transfusion should be based on reliable coagulation studies. In patients undergoing major non-cardiac surgery there is no established way of monitoring coagulation in real time. In case of intraoperative bleeding, decisions to transfuse fresh frozen plasma (FFP) and procoagulatory drugs have to be taken clinically. While hemoglobin levels and platelet counts are available within 15 minutes, standard coagulation studies require one hour. Therefore, the decision to administer FFP has to be made in the absence of any data.

Point of care testing of prothrombin time ensures that one major parameter of coagulation is available in the operation theatre within reasonable time. Several point-of-care tests for standard coagulation parameters and two different systems for thromboelastography are currently marketed. None of the PoC tests, however, is approved for intra-operative real time measurement of coagulation. One potential intra-operative bedside coagulation study system is thromboelastography. It is a viscoelastic test of whole blood clotting, which provides information about the initiation of coagulation, about clot firmness and clot lyses. It is costly and needs trained personnel with a workload of about 15 minutes to get first results. Its use is suggested for cardiac surgery and liver transplantation. During major surgery the use of thromboelastography is limited by costs, workload, The need for trained personnel, analysis time, and difficulties in interpreting results.

PoC testing of INR by various portable hand held devices is the other solution to this problem. Point of care testing of prothrombin time ensures that one major parameter of coagulation is available in the operation theatre within minutes. It is fast, easy to perform, inexpensive and may enable physicians to rationally determine the need for FFP.

Coagu Chek XS Plus[®] Roche Diagnostics and Alere IN Ratio 2 are two of such devices. Efficacy and accuracy of both have been proved in various study and give INR values on just a drop of blood within seconds.

However no study has evaluated the impact of rapid assessment of INR on the need for FFP transfusion and intraoperative blood loss. It is felt that usage of INR guided

vis a vis clinician discretion based component replacement therapy will optimize the use of FFP and may even result in less blood loss during surgery.

Materials and Methods

This study was conducted in surgical patients in a large tertiary care centre. Ethical clearance was taken from the local ethics committee at the proposal stage itself. Patients consent was taken after providing all necessary information prior to surgery. This study was conducted on 200 patients. The groups were randomised to two groups of Gp 1-study (S)-100 patients (point of care based transfusion management). Gp 2- control (C) - 100 patients (physician discretion based management).

Inclusion criteria: Age 18- 80yrs, ASA I, II and III. All patients listed for major surgery (any type excluding cardiac and liver transplant) - only those patients were included in whom blood loss was expected to be more than 20% of their blood volume. Opinion of treating surgeon and anaesthesiologist was considered final in this matter.

Exclusion criteria: Age <18 or >80 yrs, ASA IV or V. Patients on anti-platelet or anti-coagulant medication in last 5 days, Platelet count (preoperatively) <80,000/ml, INR (preoperatively) >1.5 and Hb (preoperatively) <9 gm%.

After premedication with intravenous morphine (0.05- 0.1 mg/kg body weight), Glycopyrrolate and ondansetron, General anaesthesia was induced with thiopentone sodium. Endotracheal intubation achieved after vecuronium. Anaesthesia was maintained using low flow nitrous oxide: oxygen mixture (fresh gas flow of 1ltr each) and Isoflurane (1 MAC) via a closed circuit cycle absorber system and mechanical ventilation with 5 – 10 ml/kg. Standard monitoring will include HR, ECG (Two lead), blood pressure (NIBP/ IABP), SpO₂, naso pharyngeal temperature. Patient warming was done with warm air blower (with a target temp above 36⁰C).

A baseline blood sample was taken for both the patient group after insertion of IV line prior to induction. This sample was tested for CBC and PT /INR. Those with Hb < 9 gm %, platelets < 80,000/ ml or INR > 1.5 were excluded from this study.

In the control group physician discretion was used to guide Blood platelet or FFP transfusion based on the institutional protocol. These are blood if Hb <8 gm %, platelets if <50,000 or between 50,000-80,000 with ongoing blood loss and FFP if bleeding >20% or >2 units blood given and repeated if physician desires. However these are guidelines and physician discretion based on clinical judgement is freely allowed.

In the test group if bleeding is estimated to be more than 20% CBC and POC INR is done. If Hb < 8 gm % blood is transfused, if platelets <50,000 then 1 Single Donor Platelets or 6 Random Donor Platelets are transfused. 2 units of FFP are transfused if POC INR >1.8. Test was repeated after 30mins and 2 more units of FFP was given if INR >1.5. Apart from demographic data all the test data were collected. At the end of surgery CBC was recorded. Total blood and blood product used were noted.

Observations and Results: The following observations were recorded in 200 patients, divided in 2 Groups: Study (POC based FFP transfusion group) and Control (physician

discretion based FFP transfusion group)

Demographic data

Age: The mean age of control group was 23.9 (± 11.9) months compared to 32.2 (± 20.6) months for the test group. There is no difference in age distribution between the groups P value is 0.3711.

Weight: The mean weight of the test group was 10.23 (± 2.594) kg compared to a mean weight of 11.815 (± 3.239) kg for the control group. The two-tailed P value is 0.0955, considered not significant. Thus the medians of the two groups did not differ significantly.


Table 1: Demographic data

Parameter	Test age	Control age
Mean	23.900	32.500
Std deviation	11.982	20.610
Minimum	7.000	11.000
Maximum	60.000	84.000
Median	20.500	30.000

Table 2: Weight

Parameter	wt test	wt control	Difference
Mean	10.230	11.815	-1.585
Std deviation	2.594	3.239	3.907
Minimum	5.000	7.000	-8.200
Maximum	15.000	18.000	3.500
Median	9.900	11.500	-2.000

Fig 1: Weight



Sex distribution: There were 9 Males in test group as compared to 11 in control group and 9 females in test group as compared to 11 in control group. The one-tailed P value is 0.5000, considered not significant. Thus both groups were comparable with respect to sex distribution.

a) VSD and associated anomalies.
 b) Tetralogy of Fallot (TOF) The test group had 15 VSD cases compared to 11 in the control group. The test group had 5 TOF cases compared 9 in the control group. The one-tailed P value is 0.6667, considered not significant.

Table 3: Sex Distribution

	Test	Control
Male	9	11
Female	11	9

Table 4: Diagnosis of the patients

Types of Surgeries	Test	Control
VSD	15	11
TOF	5	9

Diagnosis of the patients

Two types of cardiac lesions requiring surgeries involving cardio pulmonary bypass were part of the study.

Patient characteristics median (interquartile range). There were no significant differences among groups in any characteristic.

Table 5: Patient characteristics

Characteristic	LAG group (n =51)	POC group (n =51)	CD group (n =108)
Age (yr)	62 (57–70)	66 (58–71)	67 (60–72)
Male: number (%)	39 (76)	41 (80)	85 (78)
Female: number (%)	12 (24)	10 (20)	24 (22)
Weight (kg)	78 (72–88)	83 (76–95)	79 (70–87)
Height (cm)	172 (167–175)	170 (165–176)	170 (165–176)
Preoperative haemoglobin (g dl ⁻¹)	13.6 (13.1–14.2)	13.4 (12.7–14.3)	13.8 (12.9–14.7)
Preoperative platelet count ($\times 10^9$ litre ⁻¹)	248 (217–282)	218 (193–258)	230 (199–274)
Preoperative INR	0.92 (0.88–0.93)	0.93 (0.9–0.94)	0.94 (0.91–0.95)
Preoperative APTT ratio	0.92 (0.87–0.96)	0.9 (0.86–0.95)	0.94 (0.9–0.97)
Cardiopulmonary bypass time (min)	76 (66–84)	81 (69–93)	81 (70–96)

LAG = laboratory algorithm-guided, POC=point of care, CD=clinician discretion, INR=international normalized ratio

Table 6: T- test

	Group	N	Mean	Std. Deviation	Std. Error Mean
Age	1	100	46.88	16.626	1.663
	2	100	46.90	16.095	1.609
Wt	1	100	62.73	11.428	1.143
	2	100	62.06	11.670	1.167
BL Hb	1	100	11.679000	1.8774201	.1877420
	2	100	11.604000	1.8108495	.1810849
PCV	1	100	35.037000	5.6322602	.5632260
	2	100	34.812000	5.4325484	.5432548
BL PLT	1	100	140.74	33.146	3.315
	2	100	133.03	34.078	3.408
BLD loss	1	100	1060.89	355.721	35.572
	2	100	1038.40	369.189	36.919

WT: weight, BL HB: blood haemoglobin, PCV: Packed cell volume, BL PLT: Blood platelet, BLD loss: blood loss

Table 7: Test Statistics

	BL INR	BLD TR	PLT TR	FFP TR	PO HB	PO PLT
Mann-Whitney U	4690.500	4431.000	4905.000	4620.000	4736.000	4349.500
Wilcoxon W	9740.500	9481.000	9955.000	9670.000	9786.000	9399.500
Z	-.774	-1.438	-.340	-1.030	-.646	-1.590
Asymp. Sig. (2-tailed)	.439	.151	.734	.303	.518	.112

BL INR: Blood international normalised ratio, BLD TR: blood transfusion, PLT TR: platelet transfusion, FFP TR: fresh frozen plasma transfusion, PO HB: post-operative haemoglobin, PO PLT: post-operative platelet

User friendly

The Vigileo monitor and the Peditasat were easy to learn and the learning curve was very short. The use of the monitoring tool was easily explainable to the nursing and paramedical staff, who were able to use it effectively without any problem.

Hours of post-operative ventilation required for extubation

The mean duration of post-operative ventilation required for the patient to be extubated with stable haemodynamic parameters in the test group was 48.55 hrs compared to 78.4 hrs for the control group?

The maximum duration of post-operative ventilation in test group was 212 hrs, compared to 281 hrs for the control group. While the minimum duration was comparable being 18 hrs for test group and 19 hrs for control group. One-Sample Kolmogorov-Smirnov Test was done in each group to ensure normal distribution due to wide distribution of data.

Discussion

Bleeding is a frequent complication during surgery [1] usually more complex the surgery more is the bleed. Consequences of bleeding is exaggerated in moribund and anaemic patients. Surgery, especially complex surgeries are often associated with massive haemorrhage. Although there is no universal definition of massive haemorrhage, Hellstern [2] P *et al.* has defined it as follows: (i) blood loss exceeding circulating blood volume within a 24-hour period, (ii) blood loss of 50% of circulating blood volume within a 3-hour period, (iii) blood loss exceeding 150 ml/min, or (iv) blood loss that necessitates plasma and platelet transfusion.

The intraoperative administration of blood products, including packed red blood cells, platelets and fresh frozen plasma (FFP) is often life saving. However blood products

are not without complications. Complications of blood transfusions contribute considerably to perioperative morbidity [3] costs [4] and blood product resources are limited. Massive Hemorrhage is often associated with coagulopathy. Hemorrhage induces loss and consumption of coagulation factors. Once coagulopathy develops, it increases blood loss. Because coagulopathy and blood loss make a vicious circle, prevention of coagulopathy is important.

Large or massive blood transfusion rapidly may cause bleeding due to coagulation disorders. Stored packed red blood cells (PRBC) is deficient in fibrinogen, platelets, factor V and factor VIII. Large transfusions of stored blood leads to dilution of these elements [3]. A general indication to transfuse fresh frozen plasma (FFP) is blood loss of more than one circulating blood volume or PT-INR greater than 2.0 [5]. However, in massive Hemorrhage, British guidelines recommend that a therapeutic target of PT-INR should be less than 1.5, and that FFP should be given without PT-INR examination when blood loss is anticipated to exceed one circulating blood volume [6]. In such an emergency, we should also consider that thawing fresh frozen plasma will take about 30 min. Deficiency of fibrinogen develops earlier than that of any other coagulation factors: 100 mg/dl after loss of 120-170% of blood volume [2]. Since 5 units of FFP (400 ml) is equivalent to 1 g of fibrinogen, it increases the concentration by about 30 mg/dl in a person weighing 60 kg (circulating plasma volume: 3 L).

Fibrinogen concentrate or cryoprecipitate is preferable when urgent treatment of coagulopathy is required, but volume loading is not desirable. Hence it is often suggested that one or two units of fresh frozen plasma be transfused with every four units of stored blood if volume overload is not detrimental.

However not all patients receiving massive transfusion bleed Mannucci *et al.* showed that transfusion of FFP to counter hem dilution due to packed cell transfusion doesn't help. Thrombocytopenia is more likely the cause and not factor V or VIII deficiency in those cases that do bleed. [8]

Toulon *et al.* suggested that the Coagu Chek ProDM (a poc INR/PT/PTT device) allows an accurate measurement of PT

in patients undergoing haemorrhagic surgical procedures.⁹ However, the results were not identical to that obtained from the central laboratory, suggesting that the transfusion algorithms would have to be adapted accordingly. One of the main advantages gained using POC testing in their study is the ability to obtain results more rapidly. Actually, the turn-around time, defined as the elapsed time from blood sampling until availability of the results for the clinicians, was highly significantly shorter for the POC system than for the central laboratory (below 5 min vs. 60 min median value, range: 40 -> 120 min). Finally, the clinical interest of such a point-of-care monitoring of coagulation deserves to be prospectively investigated, especially in connection with the amount of transfused fresh frozen plasma units.

At present POC-OP trial is going on to determine the effectiveness of point of care prothrombin time testing to reduce the administration of FFP. It is a patient and assessor blind, single center randomized controlled parallel group trial in 220 patients aged between 18 and 90 years undergoing major surgery (any type, except cardiac surgery and liver transplantation) with an estimated blood loss during surgery exceeding 20% of the calculated total blood volume or a requirement of FFP according to the judgment of the physicians in charge.

Patients are randomized to usual care plus point of care prothrombin time testing or usual care alone without point of care testing. The primary outcome will be the relative risk to receive any FFP preoperatively. The authors state that inclusion of 110 patients per group will yield more than 80% power to detect a clinically relevant relative risk of 0.60 to receive FFP of the experimental as compared with the control group.

Conclusion

There was no difference in age distribution, sex weight of patients between the two groups. However the need for blood component therapy was higher in patients who were treated based on clinical discretion than compared to INR guided treatment. The mean duration of post-operative ventilation required for the patient to be extubated with stable haemodynamic parameters in the test group was 48.55 hrs compared to 78.4 hrs for the control group?

The maximum duration of post-operative ventilation in test group was 212 hrs, compared to 281 hrs for the control group. While the minimum duration was comparable being 18 hrs for test group and 19 hrs for control group. One-Sample Kolmogorov-Smirnov Test was done in each group to ensure normal distribution due to wide distribution of data.

References

1. Practice guidelines for perioperative blood transfusion and adjuvant therapies: an updated report by the American society of Anesthesiologists task force on perioperative blood transfusion and Adjuvant Therapies Anesthesiology. 2006; 105:198-208.
2. Hellstern P, Haubelt H. Indications for plasma in massive transfusion Thromb Res. 2002; 107(Suppl 1):S19-S22.
3. Jones J. Abuse of fresh frozen plasma Br Med J (Clin Res Ed). 1987; 295:287.
4. Spence RK. Surgical red blood cell transfusion practice policies, blood management practice guidelines

conference, Am J Surg. 1995; 170:3S-15S.

5. Irita K, Inada E, Yoshimura H, Warabi K, Tsuzaki K, Inaba S *et al.* Present status of preparatory measures for massive Hemorrhage and emergency blood transfusion in regional hospitals with an accredited department of anaesthesiology in 2006 Masui. 2009; 58:109-123.
6. British Committee for Standards in Haematology. Stainsby D, MacLennan S, Thomas D, Isaac J, Hamilton PJ. Guidelines on the management of massive blood loss, Br J Haematol. 2006; 135:634-641.
7. Mannucci PM, Fredrica AB, Sercha G. Hemostasis testing during massive blood replacement, A study of 172 cases Vox surg. 1982; 42:113-23.
8. Count RB, Harsh C, Simm JL, Maxwell NC, Hermsback DM, Carrico CJ. Hemostasis in massively transfused Trans fused trauma patients: Ann Surg 1979; 11:275-85.
9. Pierre A Toulon, Yves Ozier, Annick Ankri, Marie-Helene Fleron, Genevieve Leroux. Marc Samama: Monitoring of coagulation during haemorrhagic surgery by central laboratory and near-patient testing, Results of a multi center study, blood (ash annual meeting abstracts). 2004; 104:1043.