



# International Journal of Medical Anesthesiology

E-ISSN: 2664-3774

P-ISSN: 2664-3766

[www.anesthesiologypaper.com](http://www.anesthesiologypaper.com)

IJMA 2020; 3(3): 74-77

Received: 28-04-2020

Accepted: 15-07-2020

**Dr. Ranjini Basavaraj Nataraj**

Senior Resident, Department of Anaesthesiology, Subbaiah Institute of Medical Sciences, Shimoga, Karnataka, India

**Dr. Shankaranarayana P**

Professor, Department of Anaesthesiology, Yenepoya Medical College, Mangalore, Karnataka, India

**Dr. Ganapati P**

Professor, Department of Anaesthesiology, KVG Medical College, Sullia, Karnataka, India

**Corresponding Author:**

**Dr. Ranjini Basavaraj Nataraj**

Senior Resident, Department of Anaesthesiology, Subbaiah Institute of Medical Sciences, Shimoga, Karnataka, India

## Isobaric levobupivacaine and fentanyl with isobaric ropivacaine and fentanyl: Effects on heart rate and blood pressure

**Dr. Ranjini Basavaraj Nataraj, Dr. Shankaranarayana P and Dr. Ganapati P**

DOI: <https://doi.org/10.33545/26643766.2020.v3.i3b.151>

### Abstract

Most local anaesthetics block the unmyelinated C and myelinated A $\delta$  fibres that transmit pain impulses at the same rate. However the rate of blockade of A $\alpha$  and A $\beta$  (that carry motor impulses) depends on the physicochemical properties, pKa and lipid solubility of the individual local anaesthetic drugs. As ropivacaine is less lipid soluble when compared to bupivacaine, the blockade of A $\alpha$  and A $\beta$  is slow and hence produce less motor blockade than bupivacaine. In this study, 30 males in each group (group R and group L) satisfying the inclusion criteria. Group R: 2.6 cc of 0.75% isobaric ropivacaine (19.5mg) with 0.4cc of fentanyl (20 microgram). Group L: 2.6 cc of 0.5% isobaric levobupivacaine (13 mg) with 0.4cc of fentanyl (20 microgram). There is no statistically significant difference in systolic blood pressure between both the two groups at various intervals. There is no statistically significant difference in mean arterial pressure between the two groups. Ten patients in ropivacaine 0.75% group and seven patients in levobupivacaine 0.5% group developed hypotension which was managed by inj. mephentermine 6 mg IV.

**Keywords:** isobaric ropivacaine, heart rate, blood pressure

### Introduction

TURP is performed by inserting a resectoscope through the urethra and resecting prostatic tissue with an electrically powered cutting-coagulating metal loop. As much prostatic tissue as possible is resected, but the prostatic capsule is usually preserved. If the capsule is breached, irrigation solution (used to gently dilate the mucosal spaces, remove blood, cut tissue and debris from the operating field and enable better vision) is absorbed in larger amounts into the circulation and the periprostatic and retroperitoneal spaces [1].

TURP is associated with significant morbidity, including TUR syndrome, intraoperative and postoperative bleeding with need for blood transfusions and acute myocardial infarction. These problems must be considered along with the usual considerations, such as the general health of the patient, the length of the procedure and the patient and surgeon preferences when choosing an anaesthetic technique.

The use of regional anaesthesia for TURP can allow the anaesthesiologist the advantage of monitoring the patient's mental status intraoperatively. Excessive absorption of irrigating fluid during the procedure produces numerous problems with cardiovascular and neurologic implications. Visual disturbances, such as blurred vision and transient blindness have been reported in association with TURP. The biotransformation of absorbed glycine or ammonia has been implicated in these and other CNS abnormalities. Another potential complication during TURP is bladder perforation secondary to over distention with irrigation fluid or contact of the bladder wall with the surgeon's resectoscope [2].

The credit for introducing neuraxial block into clinical practice for the first time goes to August Bier in 1898. Local anaesthetics injected into the spinal subarachnoid space, block the nerve conduction to an extent determined by the concentration and volume injected. The sensitivity of different fibres varies by the drug employed. All types of nerve fibres are affected by the local anaesthetics, but within any one fibre type there is tendency for smaller slower conduction fibers to be more readily blocked than larger faster conducting fibres. Between fibre types however, these rules do not hold good. It is well established that

myelinated preganglionic B fibres which have a faster conduction time are about three times more sensitive to local anaesthetics than the slower unmyelinated postganglionic C fibres. Hypotension during central neural block may occur by three main mechanisms: decrease in venous return, vasodilatation and decreased cardiac output. Bladder distension during central nerve block has been shown to produce hypotension inappropriate to the level of block and vagal overactivity may contribute in the unsedated patient [3,4]. When hypotension occurs after SAB, patients often do not develop reflex tachycardia, this phenomenon may result from blockade of cardio-accelerator sympathetic fibers at T1 to T4 and possibly the "reverse" of the Bainbridge reflex. Rapid infusion of blood or saline sometimes produces an increase in heart rate if the initial HR is slow. This effect was described by Bainbridge in 1915. After SAB, HR decreases as a result of decrease in right atrial filling (secondary to systemic vasodilatation), which in turn leads to a decrease in outflow from intrinsic chronotropic stretch receptors in the right atrium and great veins. In fact, severe bradycardia and even cardiac arrest have been reported after spinal anaesthesia.

Most local anaesthetics block the unmyelinated C and myelinated A $\delta$  fibres that transmit pain impulses at the same rate. However the rate of blockade of A $\alpha$  and A $\beta$  (that carry motor impulses) depends on the physicochemical properties, pKa and lipid solubility of the individual local anaesthetic drugs. As ropivacaine is less lipid soluble when compared to bupivacaine, the blockade of A $\alpha$  and A $\beta$  is slow and hence produce less motor blockade than bupivacaine [5].

Levobupivacaine exerts its pharmacological action through reversible blockade of neuronal sodium channels. Myelinated nerves are blocked through exposure at the nodes of Ranvier more readily than unmyelinated nerves; and small nerves are blocked more easily than larger ones. In general, the progression of anaesthesia is related to the diameter, myelination and conduction velocity of the affected nerve fibers.

Levobupivacaine is an interesting alternative to bupivacaine for spinal anaesthesia with similar sensory and motor characteristics and recovery like bupivacaine. The regression of motor block occurs earlier with isobaric levobupivacaine as compared with isobaric bupivacaine. Intrathecal administration of 15mg of isobaric levobupivacaine provides an adequate sensory and motor block lasting for approximately 6.5 hrs. Smaller doses (i.e., 5-10 mg) are used in day-care surgeries [6]. At low concentrations, levobupivacaine produces a differential neuraxial block with preservation of motor function, which may be favourable for ambulatory surgery. Minimum effective local anaesthetic dose of levobupivacaine as recommended by an up and down sequential design study is 11.7mg.

## Methodology

**Study design:** Randomized clinical trial

**Study subjects:** Patients hospitalized for TURP.

30 males in each group (group R and group L) satisfying the inclusion criteria.

**Group R:** 2.6 cc of 0.75% isobaric ropivacaine (19.5mg) with 0.4cc of fentanyl (20 microgram).

**Group L:** 2.6 cc of 0.5% isobaric levobupivacaine (13 mg) with 0.4cc of fentanyl (20 microgram).

**Sample size:** A pilot study was done before starting the

actual study. From the pilot study, effect size of 0.868 was obtained. Considering alpha to be 0.05 and beta to be 0.20, the sample size was calculated. Thus the number of subjects in each group was found to be 30.

**Sampling method:** Simple random sampling

### Inclusion criteria

- Male patients aged 40 to 80 years with ASA grade I-III, scheduled for elective TURP.

### Exclusion criteria

- Patient's refusal, known case of hypersensitivity to amide group of local anaesthetics.
- Patients with medical complications like: (a) Hypertension, IHD, Valvular diseases (b) Anemia (c) Hypovolemia (d) Septicaemia (e) Coagulation disorders or on anticoagulant therapy.
- Local infection at the site of proposed puncture for spinal anaesthesia.
- Psychiatric disorders.
- Height <145 centimeters, morbid obesity (BMI  $\geq$  40 kg/m<sup>2</sup>).
- Patients who were unable to understand pain scales.
- History of chronic analgesic therapy, arthrosis or severe deformity of spine, peripheral neuropathy, mental disturbance or epilepsy.

## Results

**Table 1:** Heart rate (in bpm) in both the groups over a Period of time

Time	Ropivacaine and Fentanyl group			Levobupivacaine and Fentanyl group		
	Mean	SD	N	Mean	SD	N
At 0 minutes	80.07	12.61	30	80.07	9.40	30
At 2 minutes	81.07	11.61	30	82	8.84	30
At 4 minutes	79.87	11.62	30	80.33	9.06	30
At 10 minutes	74.83	11.30	30	71.7	9.67	30
At 16 minutes	75.76	7.88	30	73.06	11.53	30
At 20 minutes	76.36	7.57	30	74.36	9.11	30
At 25 minutes	77.50	8.60	30	75.4	10.38	30
At 40 minutes	77.77	8.77	30	76.33	6.60	30
At 60 minutes	78	0	1	77	2.82	2

There is no statistically significant difference in the mean heart rate between the two groups at various intervals. Two patients in each group developed bradycardia which was managed by inj. atropine 0.6 mg IV.

**Table 2:** Systolic blood pressure (in mm hg) in both the groups over a period of time

Time	Ropivacaine and Fentanyl group			Levobupivacaine and Fentanyl group		
	MEAN	SD	N	MEAN	SD	N
At 0 minutes	138.96	8.15	30	141.93	8.76	30
At 2 minutes	136.17	7.03	30	139.63	8.67	30
At 4 minutes	136.80	7.79	30	140.13	10.95	30
At 10 minutes	135.40	9.02	30	139.15	8.67	30
At 16 minutes	135.86	10.94	30	138.06	8.85	30
At 20 minutes	136.53	15.66	30	138.66	10.29	30
At 25 minutes	138.73	30.07	30	139.23	15.57	30
At 40 minutes	139.22	13.07	30	139.05	13.29	30
At 60 minutes	136	0	1	137	1.41	2

There is no statistically significant difference in systolic blood pressure between both the two groups at various intervals.

**Table 3:** Diastolic blood pressure (in MM HG) in both the groups over a period of time

Time	Ropivacaine and Fentanyl group			Levobupivacaine and Fentanyl group		
	Mean	Sd	N	Mean	Sd	N
At 0 minutes	80.06	8.25	30	82.76	4.92	30
At 2 minutes	77.93	6.68	30	80.8	4.83	30
At 4 minutes	77.76	5.87	30	80.7	5.92	30
At 10 minutes	76.16	5.16	30	78.23	4.21	30
At 16 minutes	76.86	8.77	30	78.96	7.59	30
At 20 minutes	77.93	9.54	30	78.8	9.07	30
At 25 minutes	76.86	15.72	30	77.3	7.95	30
At 40 minutes	76.11	9.31	30	77.27	7.20	30
At 60 minutes	78	0	1	77	1.41	2

There is no statistically significant difference in diastolic blood pressure at various intervals between both the groups.

**Table 4:** Mean arterial pressure (in mm hg) in both the groups over a period of time

Time	Ropivacaine and Fentanyl group			Levobupivacaine and Fentanyl group		
	Mean	Sd	N	Mean	Sd	N
At 0 minutes	99.69	8.21	30	102.48	6.2	30
At 2 minutes	97.34	6.79	30	100.41	6.11	30
At 4 minutes	97.44	6.51	30	100.51	7.59	30
At 10 minutes	95.90	6.44	30	98.53	5.69	30
At 16 minutes	96.52	9.49	30	98.66	8.01	30
At 20 minutes	97.46	11.58	30	98.75	9.47	30
At 25 minutes	97.48	20.50	30	97.94	10.49	30
At 40 minutes	97.14	10.56	30	97.86	9.23	30
At 60 minutes	97.33	0	1	97	1.41	2

There is no statistically significant difference in mean arterial pressure between the two groups. Ten patients in ropivacaine 0.75% group and seven patients in levobupivacaine 0.5% group developed hypotension which was managed by inj. mephentermine 6 mg IV.

## Discussion

The visceral pain sensation from the prostate and bladder neck is transmitted by afferent parasympathetic nerve fibers derived mostly from the second and third sacral roots traveling with the pelvic splanchnic nerves. Bladder sensation is supplied by sympathetic nerves of the hypo-gastric plexus, derived from nerve roots extending inferiorly from T11 to L2. Accidental bladder perforation also is recognized easily if the spinal level is limited to T10 because the patient would experience abdominal or shoulder pain. In addition, the uncomfortable sensation of bladder distention must be considered. Taking all this into consideration, TURP under spinal anaesthesia requires a block from T10-S4.

Hence regional anaesthesia for TURP offers some advantages over GA. Although laboratory monitoring of electrolytes is useful intraoperatively, a change in mental status in a conscious patient provides an early indication of electrolyte disturbances. Another benefit of regional anaesthesia for TURP is a decreased requirement for analgesics in the immediate postoperative period compared with general anaesthesia.

Lower urinary tract symptoms suggestive of BPH are

frequently encountered in ageing men. Over the age of 40, about one quarter of men suffer from BPH [1]. The mean age of patients in our study was 64.85yrs. Over the past few decades, TURP has become the "gold standard" of surgical treatment for patients with BPH.

There has been a significant transformation in the choice of local anaesthetics for use during spinal anaesthesia. A metanalysis showed that the relative risk of developing transient neurological symptoms was about seven times higher for spinal lignocaine than other local anaesthetics. With abandonment of lignocaine for use in spinal anaesthesia, bupivacaine gained popularity and widespread clinical acceptance. However, as reports of dangerous CNS and cardiotoxicity surfaced with use of bupivacaine, the quest for invention and evaluation of newer local anaesthetic molecules with a better safety profile started. Ropivacaine hydrochloride which is a pure S (-) enantiomer of bupivacaine is one of the fruits of that research and has shown a lot of promise as effective and safe local anaesthetic [7].

A number of studies have shown the use of plain ropivacaine in the dose range of 8 to 25 mg for various surgeries like arthroplasties, cesarean sections, knee arthroscopies, endoscopic procedures like TURP, etc. But the above mentioned studies found that intrathecal injection of isobaric ropivacaine produced a sensory block of very variable extent and a proportion of patients needed general anaesthesia because of inadequate distribution of block at lower doses. The dose of 15 mg of intrathecal ropivacaine was associated with a 5% inadequate anaesthesia in lower limb surgeries [8] and 20% inadequate anaesthesia in abdominal surgeries and hence in order to produce a reliable and effective sensory level for the planned surgical intervention, uniform dose of 19.5mg (2.6 ml of 0.75% isobaric ropivacaine) was selected.

Levobupivacaine is a preferred local anaesthetic due to its longer sensory block, lower cardiac and central nervous system toxicity. Mantouvalou M *et al.* used 15mg (3cc) of levobupivacaine which provided adequate sensory and motor block for abdominal surgeries. Lee YY *et al.* concluded that 2.6ml of 0.5% levobupivacaine can be used as an alternative to 0.5% racemic bupivacaine in spinal anaesthesia.

Opioids and local anaesthetics, when administered together intrathecally, have a potent synergistic analgesic effect. Intrathecal opioids enhance and extend the period of analgesia from sub therapeutic doses of local anaesthetics without prolonging recovery. Lipophilic opioids (e.g., fentanyl and sufentanyl) are increasingly being administered intrathecally as adjuncts to local anaesthetics. They enhance spinal anaesthesia without prolonging motor recovery or discharge time [9, 10].

The drug selected for subarachnoid block in our study was 13.5mg (2.6cc) 0.5% isobaric levobupivacaine + fentanyl 20µg (0.4cc), total 3cc versus 19.5mg (2.6cc) 0.75% isobaric ropivacaine + fentanyl 20µg (0.4cc), total 3cc.

All patients in our study groups were injected the study drugs into the subarachnoid space at the L4-5 interspace with patients in left lateral position and then turned supine immediately after the block. Patients were placed in lithotomy position on confirmation of sensory block higher than T10. If this was not achieved within 5min after SAB a head down tilt of the table was given to achieve a level of sensory block higher than T10.

## Conclusion

Hemodynamics were preserved both intraoperatively and postoperatively. However there was a small percentage of patients who developed significant fall in blood pressure and bradycardia which was statistically not significant between the two groups and were easily managed without any untoward effect.

## References

1. Gauchan S, Thapa C, Prasai A, Pyakure K, Joshi I, Tulachan J. Effects of intra the cal fentanyl as an adjunct to hyper baricbupivacaine in spinal anesthesia for elective caesarean section. *Nepal Med Coll J.* 2014; 16:5-8.
2. Koltka K, Uludag E, Senturk M, Yavru A, Karadeniz M, Sengul Tetal *et al.* Comparison of equipotent doses of ropivacaine-fentanyl and bupivacaine-fentanyl in spinal anaesthesia for lower abdominal surgery. *Anaesth Intensive Care.* 2009; 37(6):923-8.
3. Akcaboy EY, Akcaboy ZN, GogusN. Low dose levobupivacaine 0.5% with fentanyl in spinal anaesthesia for transurethral resection of prostate surgery. *JRes Med Sci.* 2011; 16(1):68-73.
4. Akan B, Yagan O, Bilal B, Erdem D, Gogus N. Comparison of levobupivacaine alone and in combination with fentanyl and sufentanyl in patients undergoing transurethral resection of the prostate. *J Res Med Sci.* 2013; 18(5):378-82.
5. Gupta K, Singhal AB, Gupta PK, Sharma D, Pandey MN, Singh I *et al.* Ropivacaine: anesthetic consideration in elderly patients for transurethral resection of prostate clinicaltrial. *Anesth Essays Res.* 2013; 7(2):178-82. doi:10.4103/0259-1162.118950
6. Chaudary A, Bogra J, Singh PK, Saxena S, Chandra G, Verma R *et al.* Efficacy of spinal ropivacaine versus ropivacaine with fentanyl in transurethral resection operations. *Saudi J Anaesth.* 2014; 8(1):88-91. doi:10.4103/1658-354X.125951.
7. Cuvas O, Basar H, Yeygel A, Turkyilmaz E, Sunay MM. Spinal anesthesia for transurethral resection operations: Levobupivacaine with or without fentanyl. *Middle East J Anaesthesiol.* 2010; 20(4):547-52.
8. Lee YY, Muchhal K, Chan CK, Cheung AS. Levobupivacaine and fentanyl for spinal anaesthesia: A randomized trial. *Eur. J Anaesthesiol.* 2005; 22(12):899-903.
9. Brahmhatt NP, PrajapatiI A, Upadhyay MR. Combination of low dose isobaric levobupivacaine 0.5% in spinal anaesthesia for lower abdominal and perineal surgeries. *Int J Res Med.* 2015; 4(2):55-60.
10. Edna Zohar, Yossi Noga, Brian Fredman *et al.* Intrathecal anesthesia for elderly patients undergoing short transurethral procedures: A dose-finding study. *Anesth Analg.* 2007; 104:552-4.