



International Journal of Medical Anesthesiology

E-ISSN: 2664-3774
P-ISSN: 2664-3766
www.anesthesiologypaper.com
IJMA 2019; 2(2): 84-88
Received: 10-09-2019
Accepted: 12-10-2019

Dr. Bhavana Harwani
Department of
Anaesthesiology, Critical Care
and Pain Medicine, Post
graduate, Meenakshi Medical
College, Hospital and Research
Institute, Kanchipuram, India

**Dr. Selvakumaran
Pannirselvam**
Department of
Anaesthesiology, Critical Care
and Pain Medicine, Associate
Professor, Meenakshi Medical
College, Hospital and Research
institute, Kanchipuram, India

Dr. PS Shanmugham
Department of
Anaesthesiology, Critical Care
and Pain Medicine, Professor,
Meenakshi Medical College,
Hospital and Research
institute, Kanchipuram, India

Dr. UG Thirumaaran
Department of
Anaesthesiology, Critical Care
and Pain Medicine, Professor
and Head of the Department
(H.O.D), Meenakshi Medical
College, Hospital and Research
institute, Kanchipuram, India

Corresponding Author:
Dr. Bhavana Harwani
Department of
Anaesthesiology, Critical Care
and Pain Medicine, Post
graduate, Meenakshi Medical
College, Hospital and Research
Institute, Kanchipuram, India

Airway assessment factors as a predictive marker of difficult direct laryngoscopy: A prospective study

Dr. Bhavana Harwani, Dr. Selvakumaran Pannirselvam, Dr. PS Shanmugham and Dr. UG Thirumaaran

DOI: <https://doi.org/10.33545/26643766.2019.v2.i2b.34>

Abstract

Background and objectives: Preoperative evaluation of anatomical landmarks help identify potentially difficult laryngoscopies; however, predictive reliability is unclear. Thus, this study was undertaken to identify and compare the most reliable variables, in prediction of difficult direct laryngoscopy.

Methodology: Pre-operative assessment of ten parameters using clinical and goniometric measurement were taken and consecutively predictors of difficult intubation were identified. On the day of surgery, after premedication and induction, laryngoscopy was performed. The glottic views were graded according to the Cormack and Lehane classification. Patients of Cormack Lehane class II B and above were considered as difficult to intubate.

Results: 15.4% of the cases were identified as difficult intubation. Cormack and Lehane classification had the highest diagnostic accuracy followed by Thyromental distance, sternomental distance and Modified Mallampati classification in that order. Mandibular-hyoid distance had highest sensitivity. when all ten parameters were taken into consideration, 95.7% of cases were classified correctly. Also, it classified correctly 97.7% of easy and 84.2% difficult intubation.

Conclusion: Apart from Cormack-Lehane grading, upon comparison of all the parameters, a combination of Thyromental distance, Atlanto-occipital joint extension and Modified Mallampati classification were able to correctly classify 95.3% of difficult intubation.

Keywords: difficult laryngoscopy, predictive airway assessment, endotracheal intubation

1. Introduction

Airway management occupies pivotal importance to an anaesthesiologist. For securing the airway, the gold standard is tracheal intubation through direct laryngoscopy. To ensure a safe anaesthetic technique, diligent efforts and absolute precision is required to secure and maintain a patent airway. The prime concern for the anaesthesiologist and the foremost task is the unanticipated difficult laryngoscopy and endotracheal intubation. Difficult tracheal intubation accounts for 17% of the respiratory related injuries and results in significant morbidity and mortality. In fact, up to 28% of all anaesthesia related deaths are secondary to the inability to mask ventilate or intubate^[1]. To aid the Anaesthesiologist in identifying these patients, several preoperative airway assessment tests have been proposed^[2-6]. It was conceptualised that the visualization of larynx during intubation is not affected by one but a plethora of factors, the concept of multivariate factors came into existence⁷. Despite large scale evaluations and efforts, predicting a difficult intubation employing a myriad of measurements and observations has not proven itself to be practicable or even reliable. Thus, the search for a predictive test which can provide an anaesthesiologist with ease of applicability, reliability and precision of prediction (discriminating power) continues. Thus, we proposed a prospective model to study the usefulness of ten different airway assessment predictors before surgery. They are: Modified Mallampati classification (MMC), Atlanto-occipital joint extension, Thyromental distance (TMD), Steno-mental distance (SMD), Mandibular hyoid distance (MHD), Ratio of height to thyromental distance (RHTMD), Inter-incisor distance (IID), Upper lip bite test (ULBT) or Mandibular protrusion test (MPT) & Cormack-Lehane grading (CLG).

2. Materials and Methods

This study was conducted at Meenakshi Medical College, Hospital & Research institute between January to September 2019 on two hundred and fifty-three adult patients aged between

18-65 years, assessed under ASA I- III, requiring surgery under GA with endotracheal intubation. Institutional ethical committee clearance and written informed consent from the patients were obtained prior to the proposed surgery. Patients with following were excluded from the study; Obvious airway malformations, need for rapid sequence intubation, Pregnancy and lactating mothers, Edentulous patients, cervical spine pathology requiring specific manipulation, Patients not willing to participate in the study and patients with BMI > more than 35kg/m².

2.1. Course of Action

All patients underwent a preanesthetic assessment prior to the surgery. A routine general physical examination was done on all patients along with routine laboratory investigations, ECG and chest X-ray.

2.1.1 Assessment criteria with Abbreviations

The enrolled patients were subjected to the following assessments preoperatively: Modified Mallampati classification (MMC), Atlanto-occipital joint extension (using goniometer), Thyromental distance (TMD), Stenomental distance (SMD), Mandibular hyoid distance (MHD), Ratio of height to thyromental distance (RHTMD), Inter-incisor distance (IID), Upper lip bite test (ULBT) and or Mandibular protrusion test (MPT)

2.1.2 Predictors of Difficult Intubation

Predictors of Difficult Intubation were identified as MMC grade III, IV^[8]; IID < 3.5cms, TMD < 6.5cms^[9], SMD < 12.5^[10], MHD^[13] < 4cms, RHTMD < 23.5^[14], ULBT class 3^[15], MPT grade B & C; Atlanto-occipital joint extension grade III, IV (12-21° & <12° respectively; Normal >35°)^[11, 12]. On the day of surgery, after premedication and induction, the patients' head and neck were kept in optimal intubating position with a pillow under the occiput during intubation (sniffing position), laryngoscopy was done using appropriate sized Macintosh blade and the glottic views were graded according to a modified classification scheme with five different grades based on the Cormack-Lehane scoring system described by Yentis^[16], who proposed that grade II be differentiated into IIA (partial view of the glottis) and IIB (arytenoids or posterior vocal cords only are

visible). Intubation is rarely difficult when a grade I or IIA view is achieved; grades IIB and III are associated with a significantly higher incidence of failed intubation. A Grade IV laryngoscopic view requires an alternate method of intubation.

2.2 Methods used for Analysis

The preoperative airway assessment data and the findings during intubation were used to determine the sensitivity, specificity, positive and negative predictive values for each test. Fisher exact test, Chi square test, Independent sample T-Test and McNemar's test were used to calculate statistically significant difference in sensitivity and specificity between these tests respectively.

3. Results

Of the total, 39(15.4%) had difficult intubation at laryngoscopy grade II B, III and IV. The overall prediction for difficult intubation considering all ten parameters has sensitivity 82.1%; specificity 97.2%; and diagnostic accuracy of 94.9%. A combination of all other factors except CLG, were able to correctly classify 95.3% of difficult intubation. Amongst this CLG, TMD & MMC, when grouped together, 95.3% of cases were correctly identified.

3.1 Most significant parameters

Three most significant factors apart from CLG were TMD, MMC and Atlanto-occipital joint extension; 92.9% of original grouped cases correctly classified when these three parameters were considered. While when all ten parameters were taken into consideration, the correct classification of difficult intubation was 95.7%, which implies that there is a marginal improvement in correctly identifying difficult intubation upon addition of rest seven parameters. The difficult cases were intubated with aid of either of the following: OELM (Optimal External Laryngeal Manipulation), bougie, stylet, video-laryngoscopy or fibreoptic bronchoscopy (FOB).

3.1.1 Results of various parameters

3.1.2 Diagnostic accuracy of various parameters




Fig 1: Cormack-Lehane Grading had highest diagnostic accuracy followed by Thyro-mental distance, Sternomental distance & Modified Mallampati classification

3.1.3 Sensitivity of various parameters




Fig 2: According to the study, Mandibular-hyoid (MHD) distance had highest sensitivity; inter-incisor distance (IID) was least sensitive

3.1.4 Specificity of various parameters




Fig 3: According to the study, Inter-incisor distance (IID) had highest specificity; Mandibular-hyoid (MHD) distance had least specificity

3.1.5 Positive predictive value of various parameters




Fig 4: According to the study, Inter-incisor distance (IID) had highest positive predictive value; Ratio of height to thyromental (RHTMD) distance had least positive predictive value.

3.1.6 Sample T-Test to compare mean values between Easy and Difficult laryngoscopy (Table 1)

1. The mean TMD in easy intubation is 7.74 ± 0.77 and in difficult intubation is 6.38 ± 0.91 . These two mean values are statistically highly significant ($p < 0.001$).
2. The mean SMD in easy intubation is 14.657 ± 1.2910

- and in difficult intubation is 12.769 ± 1.5124 . These two mean values are statistically highly significant ($p < 0.001$).
3. The mean MHD in easy intubation is 5.362 ± 0.9399 and in difficult intubation is 4.513 ± 0.5559 . These two mean values are statistically highly significant ($p < 0.001$).

- The mean IID in easy intubation is $5.48 \pm .537$ and in difficult intubation is $4.90 \pm .641$. These two mean values are statistically highly significant ($p < 0.001$).
- The mean RHTMD in easy intubation is 20.69 ± 1.98 and in difficult intubation is 25.11 ± 4.47 . These two mean values are statistically highly significant ($p < 0.001$).

Table 1: Independent sample T-Test to compare mean values between Easy and Difficult laryngoscopy

Predictor	Reality	N	Mean	Std. Dev	p-value*
3. TMD	Easy	214	7.74	.772	<0.001
	Difficult	39	6.38	.907	
4. SMD	Easy	214	14.657	1.2910	<0.001
	Difficult	39	12.769	1.5124	
5. MHD	Easy	214	5.362	.9399	<0.001
	Difficult	39	4.513	.5559	
6. IID	Easy	214	5.48	.537	<0.001
	Difficult	39	4.90	.641	
7. RHTMD	Easy	214	20.69	1.98	<0.001
	Difficult	39	25.11	4.47	

* $p < 0.05$ was regarded as significant

3.1.7 Chi-Square test to compare proportions between Easy and Difficult laryngoscopy (Table 2)

- Among ATLANTO-OCCIPITAL JOINT, in 12°- 21° category 30.0% is easy intubation, 70.0% is difficult intubation; in 22°- 34° category 86.2% is easy intubation, 13.8% is difficult intubation; and in >35° category 93.2% is easy intubation, only 6.8% is difficult intubation. This indicates that smaller angle leads to difficult intubation and larger angle tends to easy intubation. These proportions are statistically highly significant ($p < 0.001$).
- Among MMC, in grade 1- 94.3% is easy intubation, 5.7% is difficult intubation; in grade 2 – 87.9% is easy intubation, 12.1% is difficult intubation; in grade 3 – 50% is easy intubation, 50% is difficult intubation and in grade 4 – 33.3% is easy intubation, 66.7% is difficult intubation. This indicates that the greater grade leads to difficult intubation and lesser grade signifies easy intubation. These proportions are statistically highly significant ($p < 0.001$).
- Among ULBT, able to perform, is 69.9% easy intubation and 31.1% difficult intubation and unable to perform is 91.8% easy intubation and 8.2% difficult intubation. This interprets that positive outcome is easy intubation and negative outcome is difficult intubation. These proportions are statistically highly significant ($p < 0.001$).
- Among MPT, grade A – 92.1% is easy intubation, 7.9% is difficult intubation; grade B – 70.4% is easy intubation, 29.6% is difficult intubation and grade C – 75% is easy intubation, 25% is difficult intubation. Thus, grade A is easy intubation in comparison to grade B and C. These proportions are statistically highly significant ($p < 0.001$).
- Among CLG, grade I – 100% is easy intubation; grade 2A is 98.2% is easy intubation, 1.8% is difficult intubation; grade 2B is 4.3% is easy intubation, 95.7% is difficult intubation; grade 3A and 3B is 100% difficult intubation. This signifies that grade 1 and 2A

are easy intubation and subsequently higher grades are predictors of difficult intubation. These proportions are statistically highly significant ($p < 0.001$).

Table 2: Chi-Square test to compare proportions between Easy and Difficult laryngoscopy

		Reality					p-value
		Easy		Difficult		Total	
		N	%	N	%	N	
1. Atlanto-Occipital Joint Extension	12° - 21°	6	30.0%	14	70.0%	20	<0.001*
	22° - 34°	112	86.2%	18	13.8%	130	
	> 35°	96	93.2%	7	6.8%	103	
	Total	214	84.6%	39	15.4%	253	
2. MMC	Grade 1	115	94.3%	7	5.7%	122	<0.001*
	Grade 2	80	87.9%	11	12.1%	91	
	Grade 3	17	50.0%	17	50.0%	34	
	Grade 4	2	33.3%	4	66.7%	6	
	Total	214	84.6%	39	15.4%	253	
8. ULBT	No	58	69.9%	25	30.1%	83	<0.001
	Yes	156	91.8%	14	8.2%	170	
	Total	214	84.6%	39	15.4%	253	
9. MPT	A	151	92.1%	13	7.9%	164	<0.001*
	B	57	70.4%	24	29.6%	81	
	C	6	75.0%	2	25.0%	8	
	Total	214	84.6%	39	15.4%	253	
10. CLG	I	159	100.0%	0	0.0%	159	<0.001*
	2A	54	98.2%	1	1.8%	55	
	2B	1	4.3%	22	95.7%	23	
	3A	0	0.0%	15	100.0%	15	
	3B	0	0.0%	1	100.0%	1	
	Total	214	84.6%	39	15.4%	253	

*Fishers exact test.

4. Conclusion

Of all factors considered in the study, Cormack-Lehane grading is the ultimate parameter for correctly diagnosing difficult intubation, having both high sensitivity and specificity. Apart from Cormack-Lehane grading, upon comparison of all the parameters, a combination of Thyromental distance, Atlanto-occipital joint extension and Modified Mallampati classification were able to correctly classify 95.3% of difficult intubation. While when all ten parameters were taken into consideration, the correct classification of difficult intubation was 95.7%, which implies that there is a marginal improvement in correctly identifying difficult intubation upon addition of rest seven parameters. Hence, Thyromental distance, Modified Mallampati classification and Atlanto-occipital joint extension are the best parameters for predicting difficult intubation.

5. Acknowledgement

I wish to express my profound gratitude to my parents and to my dear friend Aditya bahety, who was always available when I needed his advice.

6. References

- Benumof JL. Definition and incidence of difficult airway. In: Benumof JL. Editor. Airway management: Principles and practice. St Louis Mosby, 1996, 121-125 (Ch 6).
- King TA, Adams AP. Failed tracheal intubation. Br J Anaeth. 1990; 65:400-14. [2223369].
- Mathea M, Hanna LS, Aldrete JA. Preoperative indices to anticipate difficult tracheal intubation. Anesth Analg

- 1989; 68:51-7.
4. Benumof JL. Management of the difficult adult airway with special emphasis on awake tracheal intubation. *Anesthesiology*. 1991; 75:1087-110. [1824555].
 5. Rocke DA, Morrag WB, Rout CC, Louws E. Relative risk analysis of factor associated with difficult intubation in obstetric Anesthesia. *Anesthesiology*. 1992; 77:67-73. [1610011].
 6. Mc Donald JS, Copta B, Csok RI. Proposed methods for predicting difficult intubation: Prospective evaluation of 1501 patients. *Anesthesiology*. 1992; 77:A1125
 7. Bannister FB, Mac Beth RG. Direct laryngoscopy and intubation. *Lancet*. 1944; 2:651-4.
 8. Samsoun GLT, Young JRB. Difficult tracheal intubation: A retrospective study. *Anaesthesia* 1987; 42:487-490. (From Mallampati SR: Recognition of the difficult airway.
 9. Benumof JL, editor: *Airway management principles and practice*, St Louis, Mosby, 1996, 132.
 10. Patil VU, Stehling LC, Zauder HL. Predicting the difficulty of intubation utilizing an intubation guide. *Anaesthesiology*. 1983; 10:32.
 11. Savva D. Prediction of difficult tracheal intubation. *Br J Anaesth* 1994; 73:149-153.
 12. Banister FB, Mc Beth RG. Direct laryngoscopy and tracheal intubation. *Lancet*. 1964; 2:651.
 13. Bellhouse CP, Dove C. Criteria for estimating likelihood of difficulty of endotracheal intubation with the Macintosh laryngoscope. *Anaesth intensive care*. 1988; 16:329.
 14. Chou HC, Wu TL. Mandibulohyoid distance in difficult laryngoscopy. *Br J Anaesth*. 1993; 71:335-9.
 15. Schmitt HJ, Kirmse M, Radespiel-Troger M. Ratio of patient's height to thyromental distance improves prediction of difficult laryngoscopy. *Anaesth Intensive Care*. 2002; 30:763-5.
 16. Khan ZH, Kashfi A, Ebrahim Khani E. A comparison of the upper lip bite test (a simple new technique) with modified Mallampati classification in predicting difficulty in endotracheal intubation: a prospective blinded study. *Anesth Analg*. 2003; 96:595-9.
 17. Yentis SM, Lee DJ: Evaluation of an improved scoring system for the grading of direct laryngoscopy, *Anesthesia*. 1998; 53:1041-1044.
 18. Vani VV, Kamath SK, Naik LD. The palm print as a sensitive predictor of difficult laryngoscopy in diabetics: a comparison with other airway evaluation indices. *Journal of Postgraduate Medicine*. 2000; 46(2):75-9.
 19. Lee A, Fan LT, Gin T, Karmakar MK, Ngan Kee WD. A systematic review (meta-analysis) of the accuracy of the Mallampati tests to predict the difficult airway. *Anesth Analg*. 2006; 102(6):1867-78.
 20. Butler PJ, Dhara SS. Prediction of difficult laryngoscopy: an assessment of the thyromental distance and Mallampati predictive tests. *Anaesth Intensive Care*. 1992; 20(2):139-42.