



# International Journal of Medical Anesthesiology

E-ISSN: 2664-3774  
P-ISSN: 2664-3766  
[www.anesthesiologypaper.com](http://www.anesthesiologypaper.com)  
IJMA 2022; 5(4): 07-09  
Received: 05-09-2022  
Accepted: 07-10-2022

**Dr. P Veena**  
Assistant Professor, Department  
of Anaesthesiology, Sutams,  
Trivandrum, Kerala, India

**Dr. Geethakumari P**  
Head of the Department of  
Anaesthesiology, Sutams,  
Trivandrum, Kerala, India

**Dr. Nimmi Raj**  
Assistant Professor, Department  
of Anaesthesiology, Sutams,  
Trivandrum, Kerala, India

## A routine observational study to evaluate the analgesic efficacy of epidural 0.2% ropivacaine in postoperative pain relief

**Dr. P Veena, Dr. Geethakumari P and Dr. Nimmi Raj**

**DOI:** <https://doi.org/10.33545/26643766.2022.v5.i4a.363>

### Abstract

**Aim:** To evaluate the analgesic efficacy of Epidural ropivacaine 0.2% in postoperative pain relief.

**Method:** A routine data based observational study involved 50 patients of ASA1, ASA2 who received Epidural 0.2% Ropivacaine and 50 patients who received Epidural 0.125% bupivacaine postoperatively. All patients were monitored for postoperative pain by the visual analogy scale (VAS), requirement of rescue analgesia, hemodynamic parameters and adverse effects.

**Results:** Postoperative Pain scores, Incidence of hypotension and delayed motor block were comparable between the observational groups.

**Conclusion:** Ropivacaine 0.2% and bupivacaine 0.125% were equally efficient in rescue analgesic requirement, but ropivacaine had a better safety profile in terms of less hypotension and lesser motor block.

**Keywords:** Bupivacaine, epidural, ropivacaine, postoperative pain

### Introduction

Effective pain control is essential for the optimum care of patients in the intraoperative and postoperative period. Epidural anesthesia is a safe and inexpensive technique with the advantage of providing surgical anesthesia and prolonged postoperative pain relief. Effective pain management is a critical component of postoperative care and contributes to fewer postoperative complications. The current trend in postoperative pain is multimodal analgesia. Epidural opioids have been used, but the associated major side effects, such as sedation, itching, urinary retention, nausea, vomiting, and respiratory depression have limited its widespread use. Epidural analgesia can be delivered as intermittent bolus doses, continuous infusion, and patient controlled infusion. Bupivacaine has been used successfully for many years for this purpose, in concentrations ranging from 0.0625% to 0.25%. Cardiac system and central nervous system (CNS) adverse effects related to bupivacaine have led to development of relatively safer drugs such as ropivacaine and levobupivacaine. Ropivacaine is a newer long-acting amide-linked local anaesthetic agent. It is a pure S-enantiomer of propivacaine with greater differentiation between sensory and motor blocks and with a better margin of safety due to reduced toxic potential. This study is aimed at evaluating analgesic efficacy of 0.2% ropivacaine Epidural.

### Materials and Methods

After obtaining written informed consent, a routine data based observational study was conducted which involved 50 patients of ASA1, ASA2 who received Epidural 0.2% Ropivacaine and 50 patients who received Epidural 0.125% bupivacaine postoperatively. All patients were monitored for postoperative pain by the visual analogy scale (VAS), requirement of rescue analgesia, hemodynamic parameters and adverse effects.

### Inclusion Criteria

Patients of ASA grades I to II of both Sexes

### Exclusion Criteria

Patients having severe cardiorespiratory illness, coagulation disorders, chronic liver disease, chronic kidney disease, infection at the local site, and with allergies, to amide, local anaesthetics were excluded from the study.

**Corresponding Author:**  
**Dr. P Veena**  
Assistant Professor, Department  
of Anaesthesiology, Sutams,  
Trivandrum, Kerala, India

All patients were preoperatively assessed as per standard ASA guidelines/ASRA guidelines with routine laboratory blood investigations, chest X-ray, 12-lead electrocardiogram (ECG) expert specialist consultation for indicated patients. Patients were kept fasting for 8 hours for solids

**Monitoring**

Standard ASA monitors were used. All patients were continuously monitored for Heart rate (HR), Respiratory rate (RR), and oxygen saturation, Non-invasive blood pressure and ECG.

On the day of surgery, IV access was secured with two wide bore cannulae, Patients were preloaded with crystalloids prior to spinal anaesthesia. All patients received combined spinal-epidural anaesthesia under all aseptic precautions, in L3-4, L4-5 space. Epidural catheter was placed under strict asepsis by loss of resistance to air technique, hanging drop test and by meniscal level fall test in epidural catheter. Postoperatively epidural infusion was started with 0.2% ropivacaine 4-5 ml/hr and was titrated according to patients pain score. Rescue analgesia was given with IV paracetamol. Visual analog scale (VAS) pain scores were assessed and recorded every 4 hourly.

Other related adverse effects such as hypotension and delayed motor recovery were also recorded. Hypotension was managed by fluid bolus and injection me phentermine 6 mg boluses if required. Requirement of rescue analgesia (IV paracetamol/opioids) was also noted.

**Statistical analysis**

Unpaired t-test for comparison between two groups (for comparison of means between two groups, numerical data which are normally distributed).Mann-Whitney U-test for comparison between two groups (for comparison of means between two groups, numerical data which are not normally distributed).

Chi-square test (for comparison of proportions between two groups, categorical data).


**Results**

**Table 1:** Demographic data


Parameters	Bupivacaine 0.125%	Ropivacaine 0.2%	P value
Mean age (years) ± SD	62.24±8.28	62.12±8.34	0.94
Sex Male female	15 (70%) 35 (30%)	14 (69%) 36 (31%)	0.895

**Table 2:** Visual analogy score

	Bupivacaine 0.125%	Ropivacaine 0.2%	P value
Day 0	3.33	3.6	0.114
Day 1	2.11	2.32	0.075



**Fig 1:** Requirement of rescue analgesia



**Fig 2:** Requirement of rescue analgesia

**Table 3:** Requirement of rescue analgesia

	Bupivacaine 0.125%	Ropivacaine 0.2%
Patients not requiring rescue analgesia	55.1%	40.5%
Patients requiring rescue analgesia	44.9%	59.5%

**Table 4:** Incidence of hypotension, delayed motor block was much less with 0.2% Ropivacaine

Adverse effect	Bupivacaine 0.125%	Ropivacaine 0.2%	P value
hypotension	5(9.8%)	1(2.3%)	0.025
Delayed motor block	4(7.3%)	1(2.3%)	0.046

**Discussion**

Optimum pain management should start before surgery. All patients should undergo a preoperative assessment that includes a section on pain management. This allows planning of optimal pain management techniques and facilitates early discussions to help alleviate fear of postoperative pain. Discussion of postoperative pain management at preoperative assessment aims to optimize patient satisfaction and reduce adverse effects. Effective pain management is underpinned by assessment and timely response. Self-reporting subjective pain scales represent the standard of acute pain assessment, allowing patients to report pain using a unidimensional scale of numbers or words. Commonly used to evaluate pain intensity, the visual analogue scale, verbal rating scale and numerical rating scale are valid, reliable and appropriate for use in monitoring postoperative pain in patients who are able to self-report. Our study emphasises on epidural analgesia for postoperative pain relief. Postoperative epidural analgesia is usually administered via a continuous infusion to maintain a level of analgesia and to minimize the cardiovascular and respiratory effects of bolus doses of local anaesthetics and opioid respectively. We have compared the rescue analgesic requirement while using 0.2% ropivacaine when compared to 0.125% bupivacaine. Epidural analgesia can be delivered as intermittent bolus doses, continuous infusion, and patient controlled infusion. Bupivacaine has been used successfully for many years for this purpose, in concentrations ranging from 0.0625% to 0.25%. Cardiac system and central nervous system (CNS) adverse effects related to bupivacaine have led to development of relatively safer drugs such as ropivacaine and levobupivacaine. Ropivacaine is a newer long-acting amide-linked local anaesthetic agent. It is a pure S enantiomer of propivacaine with greater differentiation between sensory and motor blocks and with a better margin of safety due to reduced toxic potential

**Conclusion**

Ropivacaine 0.2% and bupivacaine 0.125% were equally efficacious in terms of VAS pain scores, rescue analgesic requirement, but ropivacaine had a better safety profile in terms of less hypotension and lesser motor block

**Conflict of Interest**

Not available

**Financial Support**

Not available

**References**

1. De Leon-Casasola OA. When it comes to outcome, we need to define what a perioperative epidural technique is. *Anesthesia and Analgesia*. 2003;96:315-318.
2. Wheatley RG, Schug SA, Watson D. Safety and efficacy of postoperative epidural analgesia, *BJA: British Journal of Anaesthesia*. 2001 Jun 1;87(1):47-61. <https://doi.org/10.1093/bja/87.1.47>
3. Loper KA, Ready LB, Downey M, *et al*. Epidural and intravenous fentanyl infusions are clinically equivalent after knee surgery. *Anesth Analg*. 1990;70:72-5.
4. Lovstad RZ, Steen PA, Forsman M. Paraplegia after thoracotomy-not caused by the epidural catheter. *Acta Anaesthesiol Scand*. 1999;43:230-2.
5. Lowson SM, Alexander JI, Black AMS, Bambridge AD. Epidural diamorphine infusions with and without 0.167% bupivacaine for postoperative analgesia. *Eur J Anaesthesiol*. 1994;11:345-52.
6. Maeda M, Kosaka Y, Kushizaki H, Yamamori Y, Hashimoto K, Saito Y. Safety of continuous epidural anesthesia during heart surgery—change of coagulation-fibrinolysis under heparinization. *Masui Jpn J Anesthesiol*. 1999;48:723-30.
7. Mahon SV, Berry PD, Jackson M, Russell GN, Pennefather SH. Thoracic epidural infusions for post-thoracotomy pain: A comparison of fentanyl-bupivacaine mixtures vs. fentanyl alone. *Anaesthesia*. 1999;54:641-6.

**How to Cite This Article**

Veena P, Geethakumari P, Nimmi R. A routine observational study to evaluate the analgesic efficacy of epidural 0.2% ropivacaine in postoperative pain relief. *International Journal of Medical Anesthesiology*. 2022;5(4):07-09.

**Creative Commons (CC) License**

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.