



International Journal of Medical Anesthesiology

E-ISSN: 2664-3774
P-ISSN: 2664-3766
www.anesthesiologypaper.com
IJMA 2023; 6(2): 04-05
Received: 05-02-2023
Accepted: 12-03-2023

Dr. Ankhila D Hamand
Postgraduate, Department of
Anaesthesiology, Lokmanya
Tilak Municipal Medical
College, Sion, Mumbai,
Maharashtra, India

Dr. Hemangi Karnik
Professor, Department of
Anaesthesiology,
Lokmanyatilak Municipal
Medical College, Sion,
Mumbai, Maharashtra, India

Anaesthesia management for aneurysm clipping surgery in patients with severe aortic stenosis with regurgitation: A case report

Dr. Ankhila D Hamand and Dr. Hemangi Karnik

DOI: <https://doi.org/10.33545/26643766.2023.v6.i2a.386>

Abstract

Coexistence of both intracranial aneurysm and aortic stenosis is quiet rare but it may complicate management of either disease. Managing it is like tug of war, but a good case manager can find that delicate balance. We recently managed patient with severe aortic stenosis presented with anterior communicating aneurysmal.

Case-52year old female with complaints of severe headache and vomiting episodes since 10 days. With known case of hypertension and hypothyroid since 4 years on irregular medications posted for aneurysmal clipping. On pre assessment check-up we found pansystolic murmur on auscultation. On investigations it revealed concentric LVH with severe AS with moderate AR maintaining 60% ejection fraction. Other routine investigation were normal. Case was successfully conducted under general anaesthesia with endotracheal tube intubation with pre induction arterial line insertion. Anaesthesia management was tailored keeping in mind hemodynamic stability throughout procedure. With all standard ASA monitoring.

Keywords: Intracranial aneurysm, severe aortic stenosis

Introduction

Aortic stenosis -Condition fixed cardiac output which leads to ischemic and arrhythmic changes. It is risk factor for perioperative morbidity and mortality. Whereas aortic regurgitation with backward flow of blood during diastole phase

Ruptured intracranial aneurysm with risk of cerebral ischemia rebleeding hypertension & increased ICP. In addition, it causes aggravation of cardiac complication e.g. arrhythmia, cardiomyopathy.

So combination of both conditions & managing case with such a different requirement it's a challenging part.

Case report

We reported a case of 52 y/F, wt-70kg, BMI 30.2 kg/m² acute onset of severe headache & vomiting episodes since 10 days. She is k/c/o hypertension & hypothyroid since 4 years on irregular medications. Hr -72 bpm, NIBP-140/88 mm of hg SpO₂ -99%.

CNS: GCS-15/15 no neurological deficit.RS: WNL

CVS: Pansystolic murmur aortic and pulmonary area (H/o multiple syncopal attacks) her 2D ECHO shows: Concentric LVH, severe AS with moderate AR LVEF 60%: (AV PG 92 mm of hg, valve area 0.8 cm²)

CT brain: Acute SAH bilateral frontoparietal lobe.

DSA ACOM artery aneurysm without any sign of vasospasm-

Laboratory routine investigation: WNL, ECG: WNL.

Discussion

AS + AR for Aneurysmal clipping

Maintenance of steady depth of anaesthesia.

Smooth induction, hemodynamic stability to avoid →Cardiac complication (Ischemia, Arrhythmia, MI) and cerebral complication (IOAR, HTN, and Raised ICP)

Corresponding Author:
Dr. Ankhila D Hamand
Postgraduate, Department of
Anaesthesiology, Lokmanya
Tilak Municipal Medical
College, Sion, Mumbai,
Maharashtra, India

Proper fluid management

All measures to be taken to maintain cerebroprotection & brain relaxation (Head elevation, avoid neck compression, glycemic control, normocapnia, iv mannitol 20%)- Early recovery & post-operative blood pressure & pain management

Conclusion

Patient with severe aortic stenosis with aortic regurgitation for aneurysm clipping surgery requires-

- Pre evaluation for severity and extent of disease,
- Proper preoperative optimization
- Case based modification of anaesthesia technique
- Vigilant monitoring to get favourable outcome

Conflict of Interest

Not available

Financial Support

Not available

References

1. Sharma R, Mehta Y, Sapra H. Severe aortic stenosis and subarachnoid hemorrhage: Anesthetic management of lethal combination. *J Anaesthesiol Clin Pharmacol* 2013;29:255-7.
2. Vahanian A, Otto CM. Risk stratification of patients with aortic stenosis. *Eur Heart J*. 2010;31:416-23.
3. Bendel S, Ruokonen E, Pölonen P, Uusaro A. Propofol causes more hypotension than etomidate in patients with severe aortic stenosis: A double-blind, randomized study comparing propofol and etomidate. *Acta Anaesthesiol Scand*. 2007;51:284-9.
4. Stokes T, Richey R, Wray D. Guideline Development Group. Prophylaxis against infective endocarditis: Summary of NICE guidance. *Heart*. 2008;94:930-1.
5. Ramsay JG. Esmolol. *Can J Anaesth*. 1991;38:155-8.
6. Christ M, Sharkova Y, Geldner G, Maisch B. Preoperative and perioperative care for patients with suspected or established aortic stenosis facing noncardiac surgery.
7. Coppadoro A, Citerio G. Subarachnoid hemorrhage: An update for the intensivist. *Minerva Anesthesiol*. 2011;77:74-84.

How to Cite This Article

Hamand AD, Karnik H. Anaesthesia management for aneurysm clipping surgery in patients with severe aortic stenosis with regurgitation: A case report. *International Journal of Medical Anesthesiology*. 2023;6(2):04-05.

Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.