



# International Journal of Medical Anesthesiology

E-ISSN: 2664-3774  
P-ISSN: 2664-3766  
[www.anesthesiologypaper.com](http://www.anesthesiologypaper.com)  
IJMA 2024; 7(4): 37-40  
Received: 06-10-2024  
Accepted: 22-10-2024

**Dr. Anju**  
Post Graduate, Department of  
Anesthesiology, Shri  
Basaveshwara Medical College  
and Hospital, Chitradurga,  
Karnataka, India

**Dr. Megha GH**  
Professor and Head,  
Department of Anesthesiology,  
Shri Basaveshwara Medical  
College and Hospital,  
Chitradurga, Karnataka, India

**Corresponding Author:**  
**Dr. Megha GH**  
Professor and Head,  
Department of Anesthesiology,  
Shri Basaveshwara Medical  
College and Hospital,  
Chitradurga, Karnataka, India

## Randomised control trial of bispectral index with propofol induction 2 mg/kg versus 1.5 mg/kg during endotracheal intubation in general anaesthesia

**Dr. Anju and Dr. Megha GH**

**DOI:** <https://doi.org/10.33545/26643766.2024.v7.i4a.510>

### Abstract

Bispectral index (BIS) values correlates well with the sedative component in depth of anaesthesia by analysing processed EEG signals. BIS guided administration of drug results in reduced induction dose & early recovery. A Randomized, double blinded study was conducted on 120 adult patient undergoing elective surgeries under general anaesthesia. Sampling done by Simple Random Sampling using computer generated table with the duration of study of 15 months i.e. from May 2023, to July 2024. Induction dose titration of Propofol while keeping the target depth of anaesthesia is possible with BIS monitoring. In our study, group A patients achieved appropriate depth of anaesthesia with lesser induction dose of intravenous Propofol administration while achieving stable hemodynamics and early recovery.

**Keywords:** Bispectral index, propofol induction, endotracheal intubation

### Introduction

Intravenous Propofol is a common induction agent in general anaesthesia; dose dependent reduction in arterial blood pressure after drug administration has been demonstrated since its clinical introduction<sup>[1]</sup>.

Optimal induction dose administration of intravenous Propofol while maintaining appropriate depth of anaesthesia, stable haemodynamics gives smooth induction and extubation.

Bispectral index (BIS) values correlates well with the sedative component in depth of anaesthesia by analysing processed EEG signals<sup>[2,3]</sup>.

BIS guided administration of drug results in reduced induction dose & early recovery.

The main purpose of our study was comparison of two induction dose regimens (2 mg/Kg versus 1.5 mg/Kg) of Propofol with respect to depth of anaesthesia during induction and intubation phases of general anaesthesia using BIS monitoring.

Individualized dose titration with BIS helps in achieving desired hypnosis level at the same time avoiding adverse effects like hemodynamic instability, delayed recovery and other complications<sup>[4]</sup>.

### Methodology

A Randomized, double blinded study was conducted on 120 adult patient undergoing elective surgeries under general anaesthesia.

Sampling done by Simple Random Sampling using computer generated table with the duration of study of 15 months i.e. from May 2023, to July 2024.

**Statistical Analysis:** Sample size calculated using OPEN EPI software with significance level of 95% power of 80% Allocation ratio of 1

Therefore a sample size of 60 in each group was considered.

### Exclusion criteria

- Patients with anticipated difficult intubation
- Pregnant and lactating mother
- Patients on psychoactive medication, psychiatric patients, known or suspected EEG

Abnormalities (e.g. epilepsy and previous brain surgeries)

- Patients with abnormal kidney function, history of alcohol and drug abuse
- Deranged liver function
- Cardiovascular abnormalities

**Study population**

**Group A:** Propofol 1.5mg/Kg i.v induction (60 patients)

**Group B:** Propofol 2 mg/Kg i.v induction (60 patients)

**Technique**

Pre anaesthetic evaluation was done a day prior to surgery. Overnight fast according to ASA nil per oral status confirmed, Procedure was explained to the patient and written informed consent was taken. Anaesthetic machine and all equipments checked and kept ready along with the crash cart. Patient was shifted on to OT table & 18 G Intravenous (IV) cannula secured. ASA standard monitor was attached. Patients were randomly assigned by computer generated random table number to receive either intravenous Propofol 1.5mg/Kg (Group A) or 2 mg/Kg (Group B). All the study vital parameters required were recorded at baseline, at an interval of 1 minute upto 10 minutes before intubation followed by each minute monitoring for 10 minutes after intubation.

Premedicated (Inj. Ondansetron 0.08 mg/Kg IV; Inj Pantoprazole 40 mg IV; Inj. Glycopyrrolate 0.005 mg/Kg IV; Inj Fentanyl 2 mcg/Kg IV) and Pre-oxygenated.

Patients were induced with injection Propofol (At a dosage of 1.5 mg/Kg and 2 mg/Kg in group A and group B respectively) by anaesthesiologist who are not involved in the study and observer was blinded for the study; the loss of response to verbal command recorded.

Inj. Succinylcholine 2 mg/Kg IV was administered to facilitate intubation and produce muscle relaxation. After receiving adequate neuromuscular relaxation, intubation done.


Anaesthesia maintained with 67% Nitrous oxide, 33% oxygen and Isoflurane 0.6%, maintenance dose of Vecuronium with a tidal volume of 8-10 ml/Kg and respiratory rate of 10-12 breaths per minute.

After surgery, the patients were given reversal and were extubated following standard extubation criteria; shifted to the recovery.


Any adverse effects related to the drug and anaesthesia were noted and attended to appropriately with necessary drugs.

**Results**


Time of induction in Group B was faster compared to Group A and recovery was faster in Group A compared to Group B. Whereas in case of duration of intubation there was no significant difference.



**Graph 1:** End Tidal Carbon Dioxide Level



**Graph 2:** Bispectral Index



Graph 3: Oxygen Saturation




Fig 4: Heart rate response

**Discussion**

Induction dose titration of Propofol while keeping the target depth of anaesthesia is possible with BIS monitoring. In our study, group A patients achieved appropriate depth of anaesthesia with lesser induction dose of intravenous Propofol administration while achieving stable hemodynamics and early recovery [5].

Kumar B S *et al.* study showed that when the Bispectral index was maintained between 40 and 60 patients were haemodynamically stable and titration of doses according to the level of consciousness and thereby preventing complications like hypotension, respiratory depression with increased dosage [6].

Arya S *et al.* found no significant difference in the induction dose of Propofol when assessed clinically (loss of verbal response) or by BIS monitoring. The changes in BIS value and haemodynamic parameters are similar in both groups. Even in our study, both the groups showed no significant changes in haemodynamics, but there was difference in BIS values [7].

Shangne *et al.* showed BIS monitoring significantly reduces the consumption of Propofol for induction of anaesthesia. Incidence of hypotension was similar in both the BIS guided group as well as sleep guided, non –BIS group; Correlated with our study result concerning reducing the induction dose of Propofol without affecting patient hemodynamics [8].

**Conclusion**

Lesser induction dose of intravenous Propofol can provide adequate depth of anaesthesia and early recovery while maintaining stable haemodynamics. This fact can be used in patients where adverse haemodynamic variation is anticipated after induction.

**Conflict of Interest**

Not available

**Financial Support**

Not available

**References**

1. Krassioukov AV, Gelb AW, Weaver LC. Action of propofol on central sympathetic mechanisms controlling blood pressure. *Can J Anaesth.* 1993 Aug;40(8):761-9.
2. Upton RN, Martinez AM, Grant C. Comparison of the sedative properties of CNS 7056, midazolam, and propofol in sheep. *Br J Anaesth.* 2009 Dec;103(6):848-57.
3. McCollum JS, Dundee JW. Comparison of induction characteristics of four intravenous anaesthetic agents. *Anaesthesia.* 1986 Oct;41(10):995-1000.
4. Claeys MA, Gepts E, Camu F. Haemodynamic changes during anaesthesia induced and maintained with

- propofol. Br J Anaesth. 1988 Jan;60(1):3-9
5. Larijani GE, Gratz I, Afshar M, Jacobi AG. Clinical Pharmacology of Propofol; An intravenous anaesthetic agent. DICP. 1989 Oct;23(10):743-9
  6. Hug CC Jr, McLeskey CH, Nahrwold ML, Roizen MF, Stanley TH, Thisted RA, *et al.* Hemodynamic effects of propofol: data from over 25,000 patients. Anesth Analg. 1993 Oct;77(4):S21-9
  7. Liu YH, Qiu DJ, Jia L, Tan JT, Kang JM, Xie T, *et al.* Depth of anesthesia measured by bispectral index and postoperative mortality: A meta-analysis of observational studies. J Clin Anesth. 2019 Sep;56:119-125
  8. Zorrilla-Vaca A, Healy RJ, Wu CL, Grant MC. Relation between bispectral index measurements of anesthetic depth and postoperative mortality: A meta-analysis of observational studies. Can J Anaesth. 2017 Jun;64(6):597-607.

**How to Cite This Article**

Anju, Megha GH. Randomised control trial of bispectral index with propofol induction 2 mg/kg versus 1.5 mg/kg during endotracheal intubation in general anaesthesia. International Journal of Medical Anesthesiology. 2024; 7(4): 37-40.

**Creative Commons (CC) License**

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.